



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/70	A1	(11) International Publication Number: WO 99/06053 (43) International Publication Date: 11 February 1999 (11.02.99)
(21) International Application Number: PCT/US98/16053 (22) International Filing Date: 29 July 1998 (29.07.98) (30) Priority Data: 60/054,064 29 July 1997 (29.07.97) US (71) Applicant (for all designated States except US): MEDCO RESEARCH, INC. [US/US]; Suite 200, 85 T.W. Alexander Drive, Research Triangle Park, NC 27709 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): BARALDI, Pier, G. [IT/IT]; Via Tulipani 73, I-66100 Ferrara (IT). (74) Agents: BAUM, Allen, R. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: N ⁶ -SUBSTITUTED-ADENOSINE-5'-URONAMIDES AS ADENOSINE RECEPTOR MODULATORS		
(57) Abstract A series of adenosine-5'-uronamide derivatives bearing N ⁶ -arylurea, alkarylurea, heteroarylurea, arylcarbonyl, alkarylcarbonyl or heteroarylcarbonyl groups which have affinity and, in some cases, selectivity for the adenosine A ₁ or A ₃ receptors are disclosed. These compounds can be used in a pharmaceutical composition to treat disorders caused by excessive activation of the A ₁ or A ₃ receptors, or can be used in a diagnostic application to determine the relative binding of other compounds to the A ₁ or A ₃ receptors.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

N⁶-Substituted-Adenosine-5'-Uronamides as Adenosine Receptor Modulators

FIELD OF THE INVENTION

5 The present invention relates to certain N⁶-substituted-adenosine-5'-uronamide derivatives and their use in the practice of medicine as compounds with activity as agonists of adenosine receptors, in particular, the adenosine A₁ and A₃ receptors.

BACKGROUND OF THE INVENTION

10 Three major classes of adenosine receptors, classified as A₁, A₂, and A₃, have been characterized pharmacologically and have been defined on the basis of cloned sequences¹⁻⁴. A₁ receptors are coupled to the inhibition of
15 adenylate cyclase through G_i proteins and have also been shown to couple to other secondary messenger systems, including inhibition or stimulation of phosphoinositol turnover and activation of ion channels. A₂ receptors are further divided into two subtypes, A_{2A} and A_{2B}, at which adenosine agonists activate adenylyate cyclase with high and low affinity, respectively. The A₃
20 receptor sequence was first identified in a rat testes cDNA library, and this sequence, later cloned by homology to other G-protein coupled receptors from a rat brain cDNA library, was shown to correspond to a novel, functional adenosine receptor.

25 Many selective agonists and antagonists have been developed for the A₁¹⁰⁻¹⁵ and A_{2a}¹⁶⁻¹⁹ receptor subtypes. Some of these have shown promise as potential therapeutic agents in the treatment of hypertension¹⁷, Parkinson's disease²⁰, cognitive deficits²¹, schizophrenia²², epilepsy and renal failure²³. Selective and/or high affinity agonists and antagonists for
30 the A_{2b} receptor are not well known.

The discovery of the A₃ receptor has opened new therapeutic vistas in the purine field. In particular, the A₃ receptor mediates processes of inflammation, hypotension, and mast cell degranulation. This receptor apparently also has a role in the central nervous system. The A₃ selective agonist IB-MECA induces behavioral depression and upon chronic administration protects against cerebral ischemia. A₃ selective agonists at high concentrations were also found to induce apoptosis in HL-60 human leukemia cells. These and other findings have made the A₃ receptor a promising therapeutic target. Selective antagonists for the A₃ receptor are sought as potential antiinflammatory or possibly antiischemic agents in the brain. Recently, A₃ antagonists have been under development as antiasthmatic, antidepressant, antiarrhythmic, renal protective, antiparkinson and cognitive enhancing drugs.

Selective agonist²⁴ and antagonist²⁵⁻²⁸ ligands have been developed for the A₃ receptor. In the agonist field, 1B-MECA (N⁶-(3-iodobenzyl)adenosine-5'-methyluronamide) shows a K_i value of 1.1 nM at rat A₃ receptors and a 50-fold selectivity versus either A₁ or A_{2a} receptors²⁹. The related agonist, [125I]AB-MECA (N⁶-(4-amino-3-iodobenzyl)adenosine-5'-methyluronamide)³⁰ has become a useful radioligand for the screening of new derivatives at cloned A₃ receptors.

More recently, N⁶-(substituted phenylcarbamoyl)adenosine-5'-uronamides, where the substituent is 2-chloro, 3-chloro or 4-methoxy, have been prepared (Baraldi et al., Advance ACS Abstracts, December 15, 1995) and demonstrated affinity at A₃ receptors in the low nanomolar range (K_i values less than 10 nM). However, other closely related substituents, such as 3-bromo, showed a ten-fold loss in activity and affinity for the A₃ receptor (Baraldi et al., J. Med. Chem., 39:802-806 (1996)). Further, while the 2-chloro, 3-chloro and 4-methoxy substituents showed high affinity for the A₃ receptor and high selectivity relative to the A₂ receptor, the selectivity for the

A₁ receptor was not as high (the ratio of A₃ to A₁ activity was between 4 and 15).

5 A number of A₃ adenosine receptor agonists which have been previously synthesized are structurally related to adenosine itself, in which the ribose moiety is mainly intact. On the ribose, 5'-alkyluronamide groups are generally tolerated. Positions on the structure of adenosine providing flexibility of substitution, in general for adenosine agonists, have been the N⁶ and C₂ position. At the N⁶ position, most alkyl or aryl derivatives are A₁ selective. At the C₂ position, many C-, N-, or O-derivatives are A_{2a} selective. Benzyl derivatives at the N⁶ position have been shown to be A₃ selective.

15 It would be advantageous to provide other modulators of the A₁ and A₃ receptors, with high affinity and selectivity for these receptors with respect to the other adenosine receptors.


It is therefore an object of the present invention to provide compounds and methods of preparation and use thereof, which are agonists or partial agonists of the adenosine receptors, in particular, the adenosine A₁ and A₃ receptors.

SUMMARY OF THE INVENTION

25 Compounds useful as potent, and in some cases, selective agonists of the adenosine A₁ and A₃ receptors, and methods of preparation and use thereof, are disclosed.

30

The compounds have the following general formula:



wherein:

R is hydrogen, alkyl, substituted alkyl, or aryl;

R¹ is heteroaryl-NR-C(X)-, heteroaryl-C(X)-, alkaryl-NR-C(X)-, substituted alkaryl-NR-C(X)-, aryl-NR-C(X)-, aryl-C(X)-, alkaryl-C(X)-, or substituted alkaryl-C(X)-,

R² is hydrogen, alkyl, substituted alkyl, or aryl-NH-C(X)-, and

X is O, S, or NR, with the proviso that, when R² is H, R¹ is aryl-NH-C(X)- and X is O, the substituents on the aryl ring are not halo, methoxy or trifluoromethyl.

R³ and R⁴ are, independently selected from the group consisting of halo, ether, ester, azide, alkyl, alkoxy, carboxy, nitrile, nitro, trifluoro, aryl, alkaryl, thio, thioester, thioether, amine, amide and other substituents routinely used in the field of nucleoside chemistry to modify these positions. Such modifications are expected to provide the compounds with activity as partial agonists.

When R¹ is heteroaryl-NR-C(X)-, heteroaryl-C(X)-, aryl-C(X)-, alkaryl-C(X)-, or substituted alkaryl-C(X)-, the compounds tend to show affinity, and, in some cases, selectivity for the adenosine A₁ receptor. When R¹ is alkaryl-NR-C(X)-, substituted alkaryl-NR-C(X)-, or aryl-NR-C(X)-, the compounds tend to show affinity and selectivity for the A₃ receptor. Preferred substituents include p-sulfonamide, p-nitro, p-phenyl, 2,4-dichloro, p-methoxy, m-chloro, o-chloro and p-nitro.

The compounds can be used in a method for fully or partially inhibiting adenylate cyclase (A₁ and A₃) in a mammal, including a human. The methods involve administering an effective amount of a compound of formula I sufficient to fully or partially inhibiting adenylate cyclase in the mammal.

Additionally, the compounds can be used in competitive binding assays to determine the activity of other compounds in their ability to bind the A₁ or A₃ receptor.

The compounds can be used in a pharmaceutical formulation that includes a compound of formula I and one or more excipients. Various chemical intermediates, such as 2',3-isopropylidene-N-alkylcarboxamido adenosines, can be used to prepare the compounds.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a graph showing the effect of various adenosine agonists, 5'-(N-ethylcarboxamido)adenosine (NECA), N⁶-sulfonamido-phenylcarbonyl)adenosine-5'-N-ethyluronamide, 2-chloro-N⁶-(3-iodobenzyl)adenosine-5'-N-methyluronamide (Cl-IB-MECA), and N⁶-(4-amino-3-iodobenzyl)adenosine-5'-N-methyluronamide (I-AB-MECA) on the binding of [³⁵S]GTP-γ-S to membranes of RBL-2H3 rat mast cells.

Figures 2a and 2b are reaction schemes for preparing compounds of formula I.

DETAILED DESCRIPTION OF THE INVENTION

5

The present application discloses compounds useful as agonists or partial agonists of adenosine receptors, in particular, with activity as A₁ or A₃ agonists or partial agonists, and methods of preparation and use thereof.

10

The compounds can be used in a method for treating a mammal, including a human, with excessive activity at adenosine receptors, in particular, A₁ or A₃ receptors. The methods involve administering to the mammal an effective amount of a compound of formula I sufficient to modulate adenosine receptors in the mammal. The compounds can be used in a pharmaceutical formulation that includes a compound of formula I and one or more excipients.

15

Definitions

20

As used herein, a compound is an agonist of an adenosine A₁ or A₃ receptor if it is able to fully inhibit adenylate cyclase (A₁ and A₃) and is able to displace [¹²⁵I]-AB-MECA in a competitive binding assay.

25

As used herein, a compound is a partial agonist of an adenosine A₁ or A₃ receptor if it is able to partially inhibit adenylate cyclase (A₁ and A₃) and is able to displace [¹²⁵I]-AB-MECA in a competitive binding assay.

30

As used herein, a compound is an antagonist of an adenosine A₁ or A₃ receptor if it is able to prevent the inhibition due to an agonist and is able to displace [¹²⁵I]-AB-MECA in a competitive binding assay.

As used herein, a compound is selective for the A_1 receptor if the ratio of A_2/A_1 and A_3/A_1 activity is greater than about 50, preferably between 50 and 100, and more preferably, greater than about 100. A compound is selective for the A_3 receptor if the ratio of A_1/A_3 and A_2/A_3 activity is greater than about 50, preferably between 50 and 100, and more preferably, greater than about 100.

As used herein, the term "alkyl" refers to monovalent straight, branched or cyclic alkyl groups preferably having from 1 to 20 carbon atoms, more preferably 1 to 10 carbon atoms ("lower alkyl") and most preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *n*-hexyl, and the like. The terms "alkylene" and "lower alkylene" refer to divalent radicals of the corresponding alkane. Further, as used herein, other moieties having names derived from alkanes, such as alkoxyl, alkanoyl, alkenyl, cycloalkenyl, etc when modified by "lower," have carbon chains of ten or less carbon atoms. In those cases where the minimum number of carbons are greater than one, *e.g.*, alkenyl (minimum of two carbons) and cycloalkyl, (minimum of three carbons), it is to be understood that "lower" means at least the minimum number of carbons.

As used herein, the term "substituted alkyl" refers to an alkyl group, having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino aminoacyl, aminoacyloxy, oxyacylamino, cyano, halogen, hydroxyl, keto, thioketo, carboxyl, carboxylalkyl, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, hydroxylamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl, and mono- and di-alkylamino,

mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-heteroarylamino, mono- and di-heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from alkyl, aryl, heteroaryl and heterocyclic. As used herein, other moieties having the prefix "substituted" are intended to include one or more of the substituents listed above.

As used herein, the term "alkoxy" refers to the group "alkyl-O-", where alkyl is as defined above. Preferred alkoxy groups include, by way of example, methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, *tert*-butoxy, *sec*-butoxy, *n*-pentoxy, *n*-hexoxy, 1,2-dimethylbutoxy, and the like.

As used herein, the term "alkenyl" refers to alkenyl groups preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkenyl unsaturation. Preferred alkenyl groups include ethenyl ($-\text{CH}=\text{CH}_2$), *n*-propenyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), *iso*-propenyl ($-\text{C}(\text{CH}_3)=\text{CH}_2$), and the like.

As used herein, the term "alkynyl" refers to alkynyl groups preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkynyl unsaturation.

As used herein, the term "acyl" refers to the groups alkyl-C(O)-, substituted alkyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, aryl-C(O)-, heteroaryl-C(O)- and heterocyclic-C(O)- where alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, heteroaryl and heterocyclic are as defined herein.

As used herein, the term "acylamino" refers to the group $-\text{C}(\text{O})\text{NRR}$ where each R is independently hydrogen, alkyl, substituted alkyl, aryl,

heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

As used herein, the term "aryl" refers to an unsaturated aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple condensed (fused) rings (e.g., naphthyl or anthryl). Preferred aryls include phenyl, naphthyl and the like. Unless otherwise constrained by the definition for the aryl substituent, such aryl groups can optionally be substituted with from 1 to 5 substituents and preferably 1 to 3 substituents selected from the group consisting of hydroxy, acyl, alkyl, alkoxy, alkenyl, alkynyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, amino, substituted amino, aminoacyl, acyloxy, acylamino, alkaryl, aryl, aryloxy, azido, carboxyl, carboxylalkyl, cyano, halo, nitro, heteroaryl, heteroaryloxy, heterocyclic, heterocyclooxy, aminoacyloxy, oxyacylamino, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl, trihalomethyl. Preferred substituents include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy.

As used herein, the term "cycloalkyl" refers to cyclic alkyl groups of from 3 to 12 carbon atoms having a single cyclic ring or multiple condensed rings. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, and the like, or multiple ring structures such as adamantanyl, and the like.

As used herein, the terms "halo" or "halogen" refer to fluoro, chloro, bromo and iodo and preferably is either fluoro or chloro.

As used herein, the term "heteroaryl" refers to an aromatic carbocyclic group of from 1 to 15 carbon atoms and 1 to 4 heteroatoms

selected from the group consisting of oxygen, nitrogen and sulfur within at least one ring (if there is more than one ring).

5 Unless otherwise constrained by the definition for the heteroaryl substituent, such heteroaryl groups can be optionally substituted with from 1 to 5 substituents and preferably 1 to 3 substituents selected from the group consisting of hydroxy, acyl, alkyl, alkoxy, alkenyl, alkynyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, amino, substituted amino, aminoacyl, acyloxy, acylamino, alkaryl, aryl, aryloxy, 10 azido, carboxyl, carboxylalkyl, cyano, halo, nitro, heteroaryl, heteroaryloxy, heterocyclic, heterocycloxy, aminoacyloxy, oxyacylamino, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl, trihalomethyl. Preferred substituents 15 include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy. Such heteroaryl groups can have a single ring (e.g., pyridyl or furyl) or multiple condensed rings (e.g., indoliziny or benzothienyl).

20 "Heterocycle" or "heterocyclic" refers to a monovalent saturated or unsaturated group having a single ring or multiple condensed rings, from 1 to 15 carbon atoms and from 1 to 4 hetero atoms selected from the group consisting of nitrogen, sulfur and oxygen within the ring.

25 Unless otherwise constrained by the definition for the heterocyclic substituent, such heterocyclic groups can be optionally substituted with 1 to 5 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, aryl, aryloxy, halo, nitro, heteroaryl, thiol, thioalkoxy, substituted thioalkoxy, thioaryloxy, trihalomethyl, and the like. Such heterocyclic groups can have a single ring or multiple condensed rings.

30

As to any of the above groups that contain 1 or more substituents, it is understood, of course, that such groups do not contain any substitution or substitution patterns which are sterically impractical and/or synthetically non-feasible.

5

"Pharmaceutically acceptable salts" refers to pharmaceutically acceptable salts of a compound of Formula I, which salts are derived from a variety of organic and inorganic counter ions well known in the art and include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate, maleate, oxalate and the like can be used as the pharmaceutically acceptable salt.

10

15

The term "protecting group" or "blocking group" refers to any group which when bound to one or more hydroxyl, amino or carboxyl groups of the compounds (including intermediates thereof such as the aminolactams, aminolactones, etc.) prevents reactions from occurring at these groups and which protecting group can be removed by conventional chemical or enzymatic steps to reestablish the hydroxyl, amino or carboxyl group. Preferred removable amino blocking groups include conventional substituents such as t-butyloxycarbonyl (t-BOC), benzyloxycarbonyl (CBZ), and the like which can be removed by conventional conditions compatible with the nature of the product.

20

25

The following abbreviations are used herein: Abbreviations:
[¹²⁵I]AB-MECA, [¹²⁵I]N⁶-(4-amino-3-iodobenzyl)adenosine-5'-N-methyluronamide; CHO cells, Chinese hamster ovary cells; CGS 21680, 2-[4-[(2-carboxyethyl)phenyl]ethyl-amino]-5'-N-ethylcarboxamido adenosine; Cl-IB-MECA, 2-chloro-N⁶-(3-iodobenzyl)adenosine-5'-N-methyluronamide; (R)-PIA, (R)-N⁶-(phenylisopropyl)adenosine; DMSO, dimethylsulfoxide;

30

EDTA, ethylenediamine tetraacetic acid, I-AB-MECA, N⁶-(4-amino-3-iodobenzyl)adenosine-5'-N-methyluronamide; IB-MECA, N⁶-(3-iodobenzyl)adenosine-5'-N-methyluronamide; K_i, equilibrium inhibition constant; NECA, 5'-N-ethylcarboxamido adenosine; THF, tetrahydrofuran; Tris, tris(hydroxymethyl)aminomethane.

Compound Preparation

Those skilled in the art of organic chemistry will appreciate that reactive and fragile functional groups often must be protected prior to a particular reaction, or sequence of reactions, and then restored to their original forms after the last reaction is completed. Usually groups are protected by converting them to a relatively stable derivative. For example, a hydroxyl group may be converted to an ether group and an amine group converted to an amide or carbamate. Methods of protecting and de-protecting, also known as "blocking" and "de-blocking," are well known and widely practiced in the art, *e.g.*, see T. Green, *Protective Groups in Organic Synthesis*, John Wiley, New York (1981) or *Protective Groups in Organic Chemistry*, Ed. J.F.W. McOmie, Plenum Press, London (1973).

The compounds can be prepared as described below. Generally, the 2'- and 3'-hydroxy groups on an N-alkyl carboxamido adenosine, such as 5'-(N-ethylcarboxamido)adenosine (NECA), are protected. A suitable protecting group is an isopropylidene ring, although other protecting groups can be used and are intended to be within the scope of the invention.

2',3'-O-isopropylidene-N⁶-(substituted-carbonylamino)-adenosine-5'-alkyluronamides can be prepared from the isopropylidene protected N-alkylcarboxamido adenosines by reaction of the N⁶-amine group with an appropriate acid chloride or other suitable activated carboxylic acid derivative, such as an anhydride, using routine amidation conditions (as

shown in Figure 2a). 2',3'-O-isopropylidene-N⁶-(substituted-carbamoylamino)-adenosine-5'-alkyluronamides can be prepared from the isopropylidene protected N-alkylcarboxamido adenosines by reaction of the N⁶-amine group with an appropriate isocyanate using known chemistry (as shown in Figure 2b). The protecting groups are then removed to provide the desired compounds.

In those embodiments in which the 2' and/or 3'-hydroxy groups have been replaced with halo, ether, ester, azide, alkyl, alkoxy, carboxy, nitrile, nitro, trifluoro, aryl, alkaryl, thio, thioester, thioether, amine, or amide groups, the chemistry proceeds in substantially the same manner, except that thio and amine groups, which would interfere with the coupling chemistry, must first be protected with suitable protecting groups prior to coupling the N⁶ amine group with an acid halide or isocyanate.

15

Methods of Using the Compounds

The compounds can be used for all indications for which agonists and antagonists of the A₁ or A₃ receptor are effective.

Compounds which effectively modulate the A₁ receptor can be used for:

- Protection against hypoxia and/or ischemia induced injuries (e.g., stroke, infarction);
- Treatment of adenosine-sensitive cardiac arrhythmias;
- antinociception (i.e., analgesics);
- anticonvulsants;
- cardioprotection, short term (e.g., prior to percutaneous angioplasty (PTDA), angioplasty, and cardiac surgeries) and long term (prevention of myocardial infarction, especially in high risk patients, reduction of infarct damage, especially in high risk patients);

30

- neuroprotection: stroke prevention, stroke treatment, and the treatment of epilepsy;
- pain management generally, including different forms of neuropathic pain, e.g., diabetic neuropathy, post herpetic neuralgia;
- antilipid uses: reduction of free fatty acids, triglycerides, glucose;
- adjunct therapy in diabetes, including insulin and non-insulin dependent diabetes mellitus: stimulation of insulin secretion from the pancreas, increase in tissue sensitivity to insulin;
- treatment of GI disorders such as diarrhea, irritable bowel disease, irritable bowel syndrome, incontinence;
- treatment of glaucoma;
- treatment of sleep apnea;
- treatment of cardiac disarrhythmias (peroxysmal supraventricular tachycardia);
- use in combination with anesthesia for post surgical pain;
- treatment of inflammation;
- diagnostic uses, for example, to determine the presence of one or more of the above described medical conditions, or in a screening assay to determine the effectiveness of other compounds for binding to the A₁ Ado receptor (i.e., through competitive inhibition as determined by various binding assays), as described in Jacobson and Van Rhee, Purinergic approaches to experimental therapy, Jacobson and Jarvis, ed., Wiley, New York, 1997, pp. 101-128; Mathot et al., Brit. J. Pharmacol., 116:1957-1964 (1995); van der Wenden et al., J. Med. Chem., 38:4000-4006 (1995); and van Calenbergh, J. Med. Chem., 40:3765-3772 (1997), the contents of which are hereby incorporated by reference.

Compounds which effectively modulate the A₃ receptor can be used for:

- treating hypertension;
- 5 • mast cell degranulation;
- antitumor agents;
- treating cardiac hypoxia; and
- protection against cerebral ischemia;

as described, for example, in Jacobson, TIPS May 1998, pp. 185-191,
10 the contents of which are hereby incorporated by reference.

The compounds can be administered via any medically acceptable means. Suitable means of administration include oral, rectal, topical or
15 parenteral (including subcutaneous, intramuscular and intravenous)
administration, although oral or parenteral administration are preferred.

The amount of the compound required to be effective as agonist or partial agonist of an adenosine receptor will, of course, vary with the individual mammal being treated and is ultimately at the discretion of the
20 medical or veterinary practitioner. The factors to be considered include the condition being treated, the route of administration, the nature of the formulation, the mammal's body weight, surface area, age and general condition, and the particular compound to be administered. However, a suitable effective dose is in the range of about 0.1 µg/kg to about 10 mg/kg
25 body weight per day, preferably in the range of about 1 mg/kg to about 3 mg/kg per day.

The total daily dose may be given as a single dose, multiple doses, *e.g.*, two to six times per day, or by intravenous infusion for a selected
30 duration. Dosages above or below the range cited above are within the scope of the present invention and may be administered to the individual patient if

desired and necessary. For example, for a 75 kg mammal, a dosage range would be about 75 mg to about 220 mg per day, and a typical dose would be about 150 mg per day. If discrete multiple doses are indicated, treatment might typically be 50 mg of a compound given 3 times per day.

5

Formulations

The compounds described above are preferably administered in a formulation including an active compound, i.e., a compound of formula I, together with an acceptable carrier for the mode of administration. Suitable pharmaceutically acceptable carriers are known to those of skill in the art.

10

The compositions can optionally include other therapeutically active ingredients such as antivirals, antitumor agents, antibacterials, anti-inflammatories, analgesics, and immunosuppressants. The carrier must be pharmaceutically acceptable in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

15

The formulations can include carriers suitable for oral, rectal, topical or parenteral (including subcutaneous, intramuscular and intravenous) administration. Preferred carriers are those suitable for oral or parenteral administration.

20

Formulations suitable for parenteral administration include sterile aqueous preparations of the active compound, which are preferably isotonic with the blood of the recipient. Such formulations may contain distilled water, 5% dextrose in distilled water or saline. Suitable formulations also include concentrated solutions or solids containing the compound of formula (I) which upon dilution with an appropriate solvent give a solution suitable for parental administration.

25

For enteral administration, the compound can be incorporated into an inert carrier in discrete units such as capsules, cachets, tablets or lozenges,

30

each containing a predetermined amount of the active compound; as a powder or granules; or a suspension or solution in an aqueous liquid or non-aqueous liquid, *e.g.*, a syrup, an elixir, an emulsion or a draught. Suitable carriers may be starches or sugars and include lubricants, flavorings, binders, and other materials of the same nature.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active compound in a free-flowing form, *e.g.*, a powder or granules, optionally mixed with accessory ingredients, *e.g.*, binders, lubricants, inert diluents, surface active or dispersing agents. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered active compound with any suitable carrier.

A syrup or suspension may be made by adding the active compound to a concentrated, aqueous solution of a sugar, *e.g.*, sucrose, to which may also be added any accessory ingredients. Such accessory ingredients may include flavoring, an agent to retard crystallization of the sugar or an agent to increase the solubility of any other ingredient, *e.g.*, as a polyhydric alcohol, for example, glycerol or sorbitol.

The compounds can also be administered locally by topical application of a solution, ointment, cream, gel, lotion or polymeric material (for example, a PluronicTM, BASF), which may be prepared by conventional methods known in the art of pharmacy. In addition to the solution, ointment, cream, gel, lotion or polymeric base and the active ingredient, such topical formulations may also contain preservatives, perfumes, and additional active pharmaceutical agents.

Formulations for rectal administration may be presented as a suppository with a conventional carrier, e.g., cocoa butter or Witepsol S55 (trademark of Dynamite Nobel Chemical, Germany), for a suppository base.

5 Alternatively, the compound may be administered in liposomes or microspheres (or microparticles). Methods for preparing liposomes and microspheres for administration to a patient are well known to those of skill in the art. U.S. Patent No. 4,789,734, the contents of which are hereby incorporated by reference, describes methods for encapsulating biological
10 materials in liposomes. Essentially, the material is dissolved in an aqueous solution, the appropriate phospholipids and lipids added, along with surfactants if required, and the material dialyzed or sonicated, as necessary. A review of known methods is provided by G. Gregoriadis, Chapter 14, "Liposomes," Drug Carriers in Biology and Medicine, pp. 287-341
15 (Academic Press, 1979). Microspheres formed of polymers or proteins are well known to those skilled in the art, and can be tailored for passage through the gastrointestinal tract directly into the blood stream. Alternatively, the compound can be incorporated and the microspheres, or composite of microspheres, implanted for slow release over a period of time ranging from
20 days to months. See, for example, U.S. Patent Nos. 4,906,474, 4,925,673 and 3,625,214, the contents of which are hereby incorporated by reference.

25 Preferred microparticles are those prepared from biodegradable polymers, such as polyglycolide, polylactide and copolymers thereof. Those of skill in the art can readily determine an appropriate carrier system depending on various factors, including the desired rate of drug release and the desired dosage.

30 The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active compound

into association with a carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier or a finely divided solid carrier and then, if necessary, shaping the product
5 into desired unit dosage form.

In addition to the aforementioned ingredients, the formulations may further include one or more optional accessory ingredient(s) utilized in the art of pharmaceutical formulations, e.g., diluents, buffers, flavoring agents,
10 binders, surface active agents, thickeners, lubricants, suspending agents, preservatives (including antioxidants) and the like.

Determination of the Degree of Activity for the Compounds

The activity of the compounds can be readily determined using no
15 more than routine experimentation using, for example, any of the following assays.

Rat A₁ and A_{2A} Adenosine Receptor Binding Assay

Membrane preparations:

20 Male Wistar rats (200-250 g) can be decapitated and the whole brain (minus brainstem, striatum and cerebellum) was dissected on ice. The brain tissues can be disrupted in a Polytron (setting 5) in 20 vols of 50 mM Tris HCl, pH 7.4. The homogenate can then be centrifuged at 48,000 g for 10 min and the pellet resuspended in Tris-HCl containing 2 IU/ml adenosine
25 deaminase, type VI (Sigma Chemical Company, St. Louis, Mo., USA). After a 30 min incubation at 37°C, the membranes can be centrifuged and the pellets stored at -70°C. Striatal tissues can be homogenized with a Polytron in 25 vol of 50 mM Tris HCl buffer containing 10 mM MgCl₂ pH 7.4. The homogenate can be centrifuged at 48,000 g for 10 min at 4°C and
30 resuspended in Tris HCl buffer containing 2 IU/ml adenosine deaminase.

After a 30 min incubation at 37°C, the membranes can be centrifuged and the pellet stored at -70°C.

Radioligand binding assays:

5 Binding of [³H]-DPCPX (1,3-dipropyl-8-cyclopentylxanthine) to rat brain membranes can be performed essentially according to the method previously described by Bruns et al., Proc. Natl. Acad. Sci. U.S.A., 77:5547-5551 (1980), the contents of which are hereby incorporated by reference. In this method, displacement experiments are performed in 0.25 ml of buffer
10 containing 1 nM [³H]-DPCPX, 100 µl of diluted membranes of rat brain (100 µg of protein/assay) and at least 6-8 different concentrations of examined compounds. Non specific binding is determined in the presence of 10 µM of CHA (N⁶cyclohexyladenosine) and this is always ≤ 10% of the total binding. The incubation time is typically around 120 min at 25°C.

15

 Bound and free radioactivity can be separated by filtering the assay mixture through Whatman GF/B glass-fiber filters, using a Brandel cell harvester (Gaithersburg, MD, USA). The incubation mixture is diluted with 3 ml of ice-cold incubation buffer, rapidly vacuum filtered and the filter is
20 washed three times with 3 ml of incubation buffer. The filter bound radioactivity can be measured by liquid scintillation spectrometry. The protein concentration can be determined using known methodology, for example, using bovine albumin as a reference standard.

25

Human cloned A₃ Adenosine Receptor Binding Assay

Receptor binding assays: Binding assays can be carried out according to the method described by Salvatore et al., Proc. Natl. Acad. Sci. U.S.A., 90:10365-10369 (1993), the contents of which are hereby incorporated by reference. In saturation studies using this method, an aliquot of membranes
30 (8 mg protein/ml) from HEK-293 cells transfected with the human recombinant A₃ adenosine receptor (Research Biochemical International,

Natick, MA, USA) are incubated with from 10 to 12 different concentrations of [125]AB-MECA ranging from 0.1 to 5 nM. Competition experiments are carried out in duplicate in a final volume of 100 μ l in test tubes containing 0.3 nM [125]AB-MECA, 50 mM Tris HCl buffer, 10 mM MgCl₂, pH 7.4 and 20 μ l of diluted membranes (12.4 mg protein/ml) and at least 6 to 8 different concentrations of examined ligands. Incubation time is typically around 60 min at 37°C.

Bound and free radioactivity are separated by filtering the assay mixture through Whatman GF/B glass-fiber filters using a Brandel cell harvester. Non-specific binding is defined as binding in the presence of 50 μ M R-PIA and can be as high as about 30%. The incubation mixture is diluted with 3 ml of ice-cold incubation buffer, rapidly vacuum filtered and the filter is washed three times with 3 ml of incubation buffer. The bound radioactivity is counted in a Beckman gamma 5500B γ counter. The protein concentration can be determined according to known methodology, for example, using bovine albumin as reference standard.

Another suitable method for performing binding studies is the method reported in Baraldi et al., J. Med. Chem., 39:802-806 (1996).

Data Analysis

Inhibitory binding constant (K_i) values were calculated from those of IC₅₀ according to Cheng & Prusoff equation (Cheng and Prusoff, Biochem. Pharmacol., 22:3099-3108 (1973), the contents of which are hereby incorporated by reference), $K_i = IC_{50}/(1 + [C^*]/K_D^*)$, where $[C^*]$ is the concentration of the radioligand and K_D^* is its dissociation constant.

A weighted non linear least-squares curve fitting program, for example, LIGAND can be used for computer analysis of saturation and

inhibition experiments. Data are typically expressed as geometric mean, with 95% or 99% confidence limits in parentheses.

Pharmacology

5 All synthesized compounds have been tested for their affinity at rat A_1 and A_{2A} and human A_3 receptors.

EXAMPLES

10 The following examples illustrate aspects of this invention but should not be construed as limitations. The symbols and conventions used in these examples are intended to be consistent with those used in the contemporary, international, chemical literature, for example, the *Journal of the American Chemical Society* ("J.Am.Chem.Soc.") and *Tetrahedron*.

Experimental Section

Chemistry: Reactions were routinely monitored by thin-layer chromatography (TLC) on silica gel (precoated F254 Merck plates) and products visualized with iodine or aqueous potassium permanganate. Infrared spectra (IR) were measured on a Perkin Elmer 257 instrument. ^1H NMR were determined in CDCl_3 or DMSO-d_6 solutions with a Bruker AC 200 Spectrometer, with peak positions given in parts per million (δ) downfield from tetramethylsilane as internal standard, and J values are given in Hz. Light petroleum ether refers to the fractions boiling at 46-60°C. Melting points were determined on a Buchi-Tuttoli instrument and are uncorrected. Chromatography was performed with Merck 60-200 mesh silica gel. All products reported showed IR and ^1H NMR spectra in agreement with the assigned structures. Organic solutions were dried over anhydrous magnesium sulfate. Elemental analyses were performed by the microanalytical laboratory of Dipartimento di Chimica, University of Ferrara, and were within 0.4% of the theoretical values for C, H and N.

15
20
25
30

Example 1: General procedure for the preparation of 2',3'-O-isopropylidene-N⁶-(substituted-carbonylamino)adenosine-5'-N-ethyluronamide

5 2',3'-isopropylidene-NECA (0.43 mmol) was dissolved in freshly distilled dioxane (4 ml) and the appropriate acid chloride (1.3 eq.) and a catalytic amount of triethylamine (2-3 drops) were added. The mixture was refluxed under argon for 15 hours. The solvent was then removed under reduced pressure and the residue was purified by flash chromatography
10 (CH₂Cl₂-EtOAc 20%) to afford the desired compounds.

Example 2: Preparation of 2',3'-O-isopropylidene-N⁶-(4-biphenyl-carbonylamino)adenosine-5'-N-ethyluronamide

15 Following the procedure outlined in Example 1, the title compound was prepared in an 80% yield as a pale yellow foam. IR (neat) cm⁻¹ 3445, 1720, 1640, 1575; ¹H NMR (CDCl₃) δ: 0.84 (t, 3H, J = 7); 1.39 (s, 3H); 1.62 (s, 3H); 3.02-3.11 (m, 2H); 4.71 (d, 1H, J = 2); 5.39-5.46 (m, 2H); 6.19 (d, 1H, J = 2); 6.66 (t, 1H, J = 2); 7.40-7.74 (m, 7H); 8.09-8.16 (m, 3H); 8.74 (s, 1H); 9.44 (bs, 1H). Anal. (C₂₈H₂₈N₆O₅) Calc'd.: C, 63.63; H, 5.34, N, 15.90. Found: C, 63.80; H, 5.37, N, 15.82.

Example 3: Preparation of 2',3'-O-isopropylidene-N⁶-(2,4-dichlorobenzyl-carbonylamino)adenosine-5'-N-ethyluronamide

25 Following the procedure outlined in Example 1, the title compound was prepared in an 67% yield as a white foam. IR (neat) cm⁻¹ 3440, 1730, 1625, 1565, 1210; ¹H NMR (CDCl₃) δ: 0.82 (t, 3H, J = 7); 1.17 (s, 3H); 1.26 (s, 3H); 2.99-3.13 (m, 2H); 4.32 (s, 2H); 4.72 (d, 1H, J = 1.8); 5.38-5.48 (m, 2H); 6.16 (d, 1H, J = 1.8); 6.56 (t, 1H, J = 2); 7.20-7.32 (m, 3H); 7.41 (s, 1H); 8.20 (s, 1H); 8.67 (s, 1H); Anal. (C₂₃H₂₄N₇O₅Cl₂) Calc'd.: C, 57.73; H, 5.06, N, 20.49. Found: C, 57.62; H, 4.99, N, 20.42.

Example 4: Preparation of 2',3'-O-isopropylidene-N⁶-(4-methoxyphenyl-carbonylamino)adenosine-5'-N-ethyluronamide

5 Following the procedure outlined in Example 1, the title compound was prepared in an 85% yield as a white solid. IR (neat) cm^{-1} 3445, 1715, 1640, 1555, 1210; ^1H NMR (CDCl_3) δ : 0.85 (t, 3H, $J = 7$); 1.40 (s, 3H); 1.63 (s, 3H); 3.03-3.12 (m, 2H); 3.90 (s, 3H); 4.72 (d, 1H, $J = 2$); 5.41-5.49 (m, 2H); 6.16 (d, 1H, $J = 2$); 6.61 (t, 1H, $J = 2$); 7.01 (d, 2H, $J = 9$); 8.02 (s, 2H, $J = 9$); 8.10 (s, 1H); 8.74 (s, 1H); 9.09 (bs, 1H). Anal. ($\text{C}_{23}\text{H}_{26}\text{N}_6\text{O}_6$) Calc'd.: C, 57.26; H, 5.43, N, 17.42. Found: C, 57.15; H, 5.35, N, 17.28.

Example 5: Preparation of 2',3'-O-isopropylidene-N⁶-(2-chlorophenyl-carbonylamino)adenosine-5'-N-ethyluronamide

15 Following the procedure outlined in Example 1, the title compound was prepared in an 72% yield as a white foam. IR (neat) cm^{-1} 3435, 1720, 1620, 1550, 1230; ^1H NMR (CDCl_3) δ : 0.83 (t, 3H, $J = 7$); 1.39 (s, 3H); 1.62 (s, 3H); 3.00-3.07 (m, 2H); 4.71 (d, 1H, $J = 2$); 5.41-5.44 (m, 2H); 6.18 (d, 1H, $J = 2$); 6.67 (t, 1H, $J = 2$); 7.30-7.44 (m, 3H); 7.71-7.76 (m, 1H); 8.16 (s, 1H); 8.69 (s, 1H); 10.24 (bs, 1H). Anal. ($\text{C}_{22}\text{H}_{23}\text{N}_7\text{O}_5\text{Cl}$) Calc'd.: C, 52.75; H, 4.63, N, 19.57. Found: C, 52.59; H, 4.58, N, 19.43.

Example 6: Preparation of 2',3'-O-isopropylidene-N⁶-(phenyl-carbonylamino)adenosine-5'-N-ethyluronamide

25 Following the procedure outlined in Example 1, the title compound was prepared in an 90% yield as a pale yellow foam. IR (neat) cm^{-1} 3425, 1730, 1640, 1555, 1240; ^1H NMR (CDCl_3) δ : 0.79 (t, 3H, $J = 7$); 1.39 (s, 3H); 1.61 (s, 3H); 2.91-3.03 (m, 2H); 4.71 (d, 1H, $J = 2$); 5.45-5.52 (m, 2H); 6.18 (d, 1H, $J = 2$); 6.40 (t, 1H, $J = 2$); 7.27-7.53 (m, 3H); 7.83-7.87 (m, 2H); 8.19 (s, 1H); 8.63 (s, 1H); 9.18 (bs, 1H). Anal. ($\text{C}_{22}\text{H}_{24}\text{N}_6\text{O}_5$) Calc'd.: C, 58.40; H, 5.35, N, 18.57. Found: C, 58.23; H, 5.28, N, 18.45.

Example 7: General procedure for the preparation of 2',3'-O-isopropylidene-N⁶-(substituted-carbamoylamino)adenosine-5'-N-ethyluronamide

5 2',3'-isopropylidene-NECA (0.43 mmol) was dissolved in freshly distilled THF (4 ml) and the appropriate isocyanate (1.3 eq.) and a catalytic amount of triethylamine (two drops) were added. The mixture was refluxed under argon for 18 hours. The solvent was then removed under reduced pressure and the residue was purified by flash chromatography (CH₂Cl₂-
10 EtOAc 20%) to afford the desired compounds.

Example 8: Preparation of 2',3'-O-isopropylidene-N⁶-(benzyl carbamoylamino)adenosine-5'-N-ethyluronamide

15 Following the procedure outlined in Example 7, the title compound was prepared in an 82% yield as a white solid, mp 109-111 °C. IR (KBr) cm⁻¹ 3440, 1740, 1610, 1550, 1210; ¹H NMR (CDCl₃) δ: 0.76 (t, 3H, J = 7); 1.21 (s, 3H); 1.63 (s, 3H); 2.96-3.03 (m, 2H); 4.64 (d, 1H, J = 6); 4.71 (d, 1H, J =
20 1.8); 5.44-5.48 (m, 2H); 6.16 (d, 1H, J = 2); 6.52 (t, 1H, J = 2); 7.26-7.38 (m, 5H); 8.14 (s, 1H); 8.45 (s, 1H); 8.46 (s, 1H), 9.79 (bs, 1H). Anal. (C₂₆H₃₁N₇O₅) Calc'd.: C, 59.87; H, 5.99, N, 18.80. Found: C, 60.00; H, 6.03, N, 18.88.

25 **Example 9: Preparation of 2',3'-O-isopropylidene-N⁶-(4-sulfonamido-phenylcarbamoylamino)adenosine-5'-N-ethyluronamide**

30 Following the procedure outlined in Example 7, the title compound was prepared in an 60% yield as a white solid, mp 207 °C. IR (KBr) cm⁻¹ 3450-3250, 1730, 1610, 1565, 1360, 1230; ¹H NMR (CDCl₃) δ: 0.57 (t, 3H, J = 7); 1.35 (s, 3H); 1.54 (s, 3H); 2.96-3.03 (m, 2H); 4.60 (d, 1H, J = 2); 5.47-5.50 (m, 2H); 6.47 (d, 1H, J = 2); 6.60 (t, 1H, J = 2); 7.26 (d, 2H, J = 9); 7.74 (d, 2H, J = 9); 8.60 (s, 1H); 8.64 (s, 1H), 9.22 (s, 1H); 10.47 (bs, 1H);

12.05 (s, 1H). Anal. ($C_{22}H_{26}N_8O_7S$) Calc'd.: C, 48.35; H, 4.79, N, 20.50. Found: C, 48.27; H, 4.79, N, 20.63.

5 **Example 10: Preparation of 2',3'-O-isopropylidene-N⁶-(4-acetyl-phenylcarbamoylamino)adenosine-5'-N-ethyluronamide**

Following the procedure outlined in Example 7, the title compound was prepared in an 77% yield as a white solid, mp 204°C. IR (KBr) cm^{-1}
10 3430, 1740, 1720, 1630, 1545, 1240; 1H NMR ($CDCl_3$) δ : 0.83 (t, 3H, J = 7); 1.42 (s, 3H); 1.64 (s, 3H); 2.61(s, 3H); 2.99-3.08 (m, 2H); 4.74 (d, 1H, J = 2); 5.47-5.55 (m, 2H); 6.21 (d, 1H, J = 2); 6.47 (t, 1H, J = 2); 7.74 (d, 2H, J = 9); 7.98 (d, 2H, J = 9); 8.21 (s, 1H); 8.63 (s, 1H), 8.67 (bs, 1H); 12.01 (s, 1H). Anal. ($C_{22}H_{26}N_8O_7$) Calc'd.: C, 51.36; H, 5.09, N, 21.78. Found: C,
15 51.35; H, 5.12, N, 21.66.

20 **Example 11: Preparation of 2',3'-O-isopropylidene-N⁶-((R)- α -phenylethyl-carbamoylamino)adenosine-5'-N-ethyluronamide**

Following the procedure outlined in Example 7, the title compound was prepared in an 65% yield as a white solid, mp 110-111°C. IR (KBr) cm^{-1}
 1 3430, 1730, 1620, 1555, 1230; 1H NMR ($CDCl_3$) δ : 0.68 (t, 3H, J = 7); 1.41 (s, 3H); 1.63 (s, 3H); 1.64 (d, 3H, J = 7); 2.87-2.96 (m, 2H); 4.71 (d,
25 1H, J = 1.8); 5.16 (m, 1H); 5.48-5.51 (m, 2H); 6.16 (d, 1H, J = 2); 6.52 (bs, 1H); 7.25-7.408 (m, 5H); 8.15 (s, 1H); 8.51 (s, 1H); 9.84 (bs, 1H), 10.94 (s, 1H). Anal. ($C_{23}H_{29}N_7O_5$) Calc'd.: C, 57.13; H, 6.05, N, 20.28. Found: C, 57.01; H, 5.99, N, 20.33.

30 **Example 12: Preparation of 2',3'-O-isopropylidene-N⁶-((S)- α -phenylethyl-carbamoylamino)adenosine-5'-N-ethyluronamide**

Following the procedure outlined in Example 7, the title compound was prepared in an 82% yield as a white solid, mp 109-111°C. IR (KBr) cm^{-1} 3445, 1725, 1615, 1560, 1230; ^1H NMR (CDCl_3) δ : 0.85 (t, 3H, $J = 7$); 1.40 (s, 3H); 1.63 (s, 3H); 1.64 (d, 3H, $J = 7$); 3.00-3.09 (m, 2H); 4.72 (d, 1H, $J = 1.8$); 5.17 (m, 1H); 5.39-5.50 (m, 2H); 6.14 (d, 1H, $J = 2$); 6.57 (bs, 1H); 7.26-7.45 (m, 5H); 8.12 (s, 1H); 8.51 (s, 1H); 9.81 (bs, 1H); 10.87 (s, 1H). Anal. ($\text{C}_{23}\text{H}_{29}\text{N}_7\text{O}_5$) Calc'd.: C, 57.13; H, 6.05, N, 20.28. Found: C, 57.25; H, 6.11, N, 20.37.

Example 13: Preparation of 2',3'-O-isopropylidene- N^6 -(5-methyl-isoxazol-3-yl-carbamoylamino)adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 7, the title compound was prepared in an 83% yield as a white solid, mp 120°C. IR (KBr) cm^{-1} 3450, 1720, 1620, 1550, 1215; ^1H NMR (CDCl_3) δ : 0.82 (t, 3H, $J = 7$); 1.20 (s, 3H); 1.41 (s, 3H); 2.44 (s, 3H); 3.00-3.07 (m, 2H); 4.74 (d, 1H, $J = 2$); 5.46-5.50 (m, 2H); 6.21 (d, 1H, $J = 2$); 6.55 (t, 1H, $J = 2$); 6.70 (s, 1H); 8.37 (s, 1H); 8.63 (s, 1H); 9.39 (bs, 1H); 12.34 (bs, 1H). Anal. ($\text{C}_{20}\text{H}_{24}\text{N}_8\text{O}_6$) Calc'd.: C, 50.84; H, 5.12, N, 23.72. Found: C, 50.96; H, 5.18, N, 23.64.

Example 14: Preparation of 2',3'-O-isopropylidene- N^6 -(1,3,4-thiadiazol-2-yl-carbamoylamino)adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 7, the title compound was prepared in a 74% yield as a yellow solid, mp 148°C. IR (KBr) cm^{-1} 3430, 1730, 1615, 1560, 1240; ^1H NMR (CDCl_3) δ : 0.86 (t, 3H, $J = 7$); 1.44 (s, 3H); 1.67 (s, 3H); 3.07-3.14 (m, 2H); 4.78 (d, 1H, $J = 1.8$); 4.78 (d, 1H, $J = 1.8$); 5.47-5.51 (m, 2H); 6.24 (d, 1H, $J = 1.8$); 6.52 (t, 1H, $J = 2$); 8.36 (s, 1H); 8.75 (s, 1H); 8.91 (bs, 1H); 9.40 (bs, 1H); 11.72 (bs, 1H). Anal. ($\text{C}_{18}\text{H}_{21}\text{N}_9\text{O}_5\text{S}$) Calc'd.: C, 45.47; H, 4.45, N, 26.51. Found: C, 45.39; H, 4.44, N, 26.46.

Example 15: Preparation of 2',3'-O-isopropylidene-N⁶-(4-n-propyloxy-phenylcarbamoylamino)adenosine-5'-N-ethyluronamide

5

Following the procedure outlined in Example 7, the title compound was prepared in an 70% yield as a pale yellow foam. IR (neat) cm^{-1} 3440, 1725, 1620, 1555, 1220; ^1H NMR (CDCl_3) δ : 0.71 (t, 3H, $J = 7$); 0.90 (s, 3H, $J = 7$); 1.40 (s, 3H); 1.62 (s, 3H); 1.81-2.03 (m, 2H); 2.95-3.01 (m, 2H); 3.90 (t, 2H, $J = 7$); 4.71 (d, 1H, $J = 1.8$); 5.46-5.51 (m, 2H); 6.19 (d, 1H, $J = 1.8$); 6.51 (t, 1H, $J = 2$); 6.89 (d, 2H, $J = 9$); 7.48 (d, 2H, $J = 9$); 8.25 (s, 1H); 8.56 (s, 1H); 8.80 (bs, 1H), 11.48 (s, 1H). Anal. ($\text{C}_{22}\text{H}_{25}\text{N}_7\text{O}_5$) Calc'd.: C, 56.52; H, 5.39, N, 20.97. Found: C, 56.44; H, 5.42, N, 21.03.

10

15

Example 16: Preparation of 2',3'-O-isopropylidene-N⁶-bis-(4-nitrophenyl-carbamoylamino)adenosine-5'-N-ethyluronamide

20

Following the procedure outlined in Example 7, the title compound was prepared in an 65% yield as a yellow solid, mp 261 °C. IR (KBr) cm^{-1} 3430, 1730, 1620, 1555, 1210; ^1H NMR ($\text{DMSO}-d_6$) δ : 0.56 (t, 3H, $J = 7$); 1.34 (s, 3H); 1.54 (s, 3H); 2.67-2.83 (m, 2H); 4.61 (d, 1H, $J = 1.8$); 5.47-5.50 (m, 2H); 6.47 (d, 1H, $J = 1.8$); 7.61 (t, 1H, $J = 2$); 7.71 (d, 2H, $J = 9$); 7.91 (d, 2H, $J = 9$); 8.22 (d, 2H, $J = 9$); 8.24 (s, 1H), 8.64 (d, 2H, $J = 9$); 9.72 (s, 1H); 10.63 (bs, 1H); 12.24 (s, 1H). Anal. ($\text{C}_{29}\text{H}_{28}\text{N}_{10}\text{O}_{10}$) Calc'd.: C, 51.48; H, 4.17, N, 20.70. Found: C, 51.56 ; H, 4.16, N, 20.69.

25

30

Example 17: Preparation of 2',3'-O-isopropylidene-N⁶-bis-(5-chloro-pyridin-2-yl-carbamoylamino)adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 7, the title compound was prepared in a 60% yield as a yellow foam. IR (neat) cm^{-1} 3440, 1710, 1630, 1540, 1220; ^1H NMR (CDCl_3) δ : 0.58 (t, 3H, $J = 7$); 1.35 (s, 3H); 1.54 (s, 3H); 2.71-2.85 (m, 2H); 4.60 (d, 1H, $J = 2$); 5.47-5.51 (m, 2H); 6.47 (d,

1H, J = 2); 7.52 (t, 1H, J = 2); 7.79-7.92 (m, 4H); 8.34-8.38 (m, 2H); 8.60 (s, 1H); 8.65 (s, 1H), 10.66 (bs, 1H); 12.22 (bs, 1H). Anal. (C₂₇H₂₆N₈O₆Cl₂) Calc'd.: C, 51.52; H, 4.16, N, 17.80. Found: C, 51.55; H, 4.13, N, 17.84.

5 **Example 18: General procedure for the preparation of N⁶-(substituted)adenosine-5'-N-ethyluronamide**

10 A solution of the isopropylidene derivative (0.084 mmol) in aqueous 1N HCl (5 ml) and dioxane (5 ml) was stirred at 65 °C for 1 hour. The solvent was then removed under reduced pressure and the residue was crystallized from ethanol to afford the desired compounds.

15 **Example 19: Preparation of N⁶-(4-biphenyl-carbonylamino)-adenosine-5'-N-ethyluronamide**

20 Following the procedure outlined in Example 18, the title compound was prepared in a 65% yield as a white solid, mp 221 °C. IR (KBr) cm⁻¹ 3500-3100, 1720, 1615, 1550; ¹H NMR (DMSO d₆) δ: 1.07 (t, 3H, J = 7); 3.17-3.24 (m, 2H); 4.23-4.25 (m, 1H); 4.36 (d, 1H, J = 2); 4.71-4.73 (m, 1H); 5.70 (d, 1H, J = 8); 5.79 (d, 1H, J = 4); 6.13 (d, 1H, J = 8); 7.43-7.53 (m, 3H); 7.78-7.89 (m, 4H); 8.16 (d, 2H, J = 11); 8.46 (t, 1H, J = 4), 8.80 (s, 1H); 8.82 (s, 1H); 11.34 (bs, 1H). Anal. (C₂₅H₂₄N₆O₅) Calc'd.: C, 61.47; H, 4.95, N, 17.20. Found: C, 61.57; H, 5.00, N, 17.28.

25 **Example 20: Preparation of N⁶-(2,4-dichlorobenzyl-carbonylamino)-adenosine-5'-N-ethyluronamide**

30 Following the procedure outlined in Example 18, the title compound was prepared in a 55% yield as a white solid, mp 132-134 °C. IR (KBr) cm⁻¹ 3450-3050, 1730, 1635, 1545, 1230; ¹H NMR (DMSO d₆) δ: 1.08 (t, 3H, J = 7); 3.18-3.25 (m, 2H); 3.34 (s, 2H); 4.12-4.15 (m, 1H); 4.30 (s, 1H); 4.49-4.62 (m, 2H); 5.57 (d, 1H, J = 8); 5.77 (d, 1H, J = 4); 5.95 (d, 1H, J = 8); 7.44-7.51 (m, 3H); 8.19 (s, 1H); 8.39 (s, 1H), 8.95 (bs, 1H). Anal.

(C₂₀H₂₀N₆O₅) Calc'd.: C, 56.60; H, 4.75, N, 19.80. Found: C, 56.60; H, 4.77, N, 19.84.

Example 21: Preparation of N⁶-(4-methoxyphenyl-carbonylamino)-adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 18, the title compound was prepared in an 80% yield as a white solid, mp 167-169°C. IR (KBr) cm⁻¹ 3550-3150, 1710, 1640, 1530, 1270; ¹H NMR (DMSO d₆) δ: 1.07 (t, 3H, J = 7); 3.15-3.23 (m, 2H); 3.86 (s, 3H); 4.20-4.24 (m, 1H); 4.35 (d, 1H, J = 2); 4.70-4.75 (m, 1H); 5.69 (d, 1H, J = 7); 5.77 (d, 1H, J = 4); 6.11 (d, 1H, J = 7); 7.08 (d, 2H, J = 9); 8.04 (d, 2H, J = 9); 8.44 (bs, 1H); 8.77 (s, 1H); 8.78 (s, 1H), 11.12 (s, 1H). Anal. (C₂₀H₂₂N₆O₆) Calc'd.: C, 54.30; H, 5.01, N, 18.99. Found: C, 54.38; H, 4.98, N, 19.02.

Example 22: Preparation of N⁶-(4-chlorophenyl-carbonylamino)-adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 18, the title compound was prepared in a 78% yield as a white solid, mp 179-180°C. IR (KBr) cm⁻¹ 3510-3050, 1720, 1660, 1510, 1250; ¹H NMR (DMSO d₆) δ: 1.06 (t, 3H, J = 7); 3.15-3.22 (m, 2H); 4.22 (bs, 1H); 4.34 (s, 1H); 4.68-4.70 (m, 1H); 5.66 (d, 1H, J = 7); 5.76 (d, 1H, J = 4); 6.10 (d, 1H, J = 7); 7.43-7.61 (m, 4H); 8.44 (bs, 1H); 8.67 (s, 1H); 8.82 (bs, 1H); 11.53 (bs, 1H). Anal. (C₁₉H₁₉N₆O₅) Calc'd.: C, 55.47; H, 4.66, N, 20.43. Found: C, 55.53; H, 4.72, N, 20.50.

Example 23: Preparation of N⁶-(phenyl-carbonylamino)-adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 18, the title compound was prepared in a 67% yield as a white solid, mp 157-159°C. IR (KBr) cm⁻¹ 3500-3000, 1715, 1630, 1535, 1260; ¹H NMR (DMSO d₆) δ: 1.03 (t, 3H, J =

7); 3.16-3.19 (m, 2H); 3.74 (bs, 1H); 4.22 (bs, 1H); 4.36 (s, 1H); 4.51-4.63 (m, 1H); 6.13 (d, 1H, J = 6.8); 7.48-7.59 (m, 3H); 8.04-8.11 (m, 2H); 8.52 (bs, 1H), 8.79 (s, 1H); 8.83 (s, 1H); 11.22 (s, 1H). Anal. (C₁₉H₂₀N₆O₅) Calc'd.: C, 55.34; H, 4.89, N, 20.38. Found: C, 55.47; H, 4.93, N, 20.43.

5

Example 24: Preparation of N⁶-(benzylcarbamoylamino)-adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 18, the title compound was prepared in a 70% yield as a white solid, mp 184-186°C. IR (KBr) cm⁻¹ 3500-3100, 1675, 1620, 1565, 1520, 1310; ¹H NMR (DMSO d₆) δ: 1.06 (t, 3H, J = 7); 3.10-3.20 (m, 2H); 4.05 (bs, 1H); 4.17-4.20 (m, 1H); 4.36 (s, 1H); 4.47-4.50 (m, 2H); 4.55-4.65 (m, 1H); 6.05-6.10 (m, 1H); 7.17-7.40 (m, 5H); 8.30-8.40 (m, 1H); 8.62 (s, 1H); 8.86 (s, 1H); 9.50-9.60 (m, 1H); 10.30 (bs, 1H). Anal. (C₂₀H₂₄N₇O₅) Calc'd.: C, 54.29; H, 5.47, N, 22.16. Found: C, 54.36; H, 5.52, N, 22.08.

15

Example 25: Preparation of N⁶-(4-sulfonamido-phenylcarbamoyl)-adenosine-5'-N-ethyluronamide

20

Following the procedure outlined in Example 18, the title compound was prepared in a 63% yield as a white solid, mp 183-185°C. IR (KBr) cm⁻¹ 3550-3050, 1715, 1630, 1545, 1370, 1250; ¹H NMR (DMSO d₆) δ: 1.07 (t, 3H, J = 7); 3.16-3.26 (m, 2H); 4.21 (bs, 1H); 4.36 (s, 1H); 4.63-4.69 (m, 1H); 5.62 (d, 1H, J = 6.8); 5.78 (d, 1H, J = 4); 6.11 (d, 1H, J = 6.8); 7.31 (bs, 2H); 7.81 (s, 4H); 8.49 (bs, 1H); 8.75 (s, 1H), 8.89 (s, 1H); 9.69 (bs, 1H); 11.92 (bs, 1H). Anal. (C₁₉H₂₂N₈O₇S) Calc'd.: C, 45.06; H, 4.38, N, 22.12. Found: C, 45.15; H, 4.36, N, 22.07.

25

30

Example 26: Preparation of N⁶-(4-acetyl-phenylcarbamoyl)-adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 18, the title compound was prepared in an 83% yield as a white solid, mp 187°C. IR (KBr) cm^{-1} 3550-3100, 1740, 1720, 1630, 1525, 1215; ^1H NMR (DMSO d_6) δ : 1.08 (t, 3H, $J = 7$); 2.55 (s, 2H); 3.16-3.23 (m, 2H); 4.18-4.20 (m, 1H); 4.21 (d, 1H, $J = 2$); 4.45 (bs, 1H); 4.64-4.69 (m, 1H); 6.11 (d, 1H, $J = 7$); 7.77 (d, 2H, $J = 9$); 7.98 (d, 2H, $J = 9$); 8.49 (bs, 1H), 8.76 (s, 1H); 8.88 (s, 1H); 10.62 (bs, 1H); 11.94 (bs, 1H). Anal. ($\text{C}_{21}\text{H}_{23}\text{N}_7\text{O}_6$) Calc'd.: C, 53.73; H, 4.94, N, 20.89. Found: C, 53.80; H, 4.95, N, 20.93.

Example 27: Preparation of N^6 -((R)- α -phenylethylcarbamoyl)-adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 18, the title compound was prepared in a 65% yield as a white solid, mp 153-155°C. IR (KBr) cm^{-1} 3550-3100, 1720, 1635, 1535, 1240; ^1H NMR (DMSO d_6) δ : 1.05 (t, 3H, $J = 7$); 1.51 (d, 3H, $J = 8$); 3.14-3.21 (m, 2H); 3.35-3.41 (m, 1H); 4.20 (s, 1H); 4.36 (s, 1H); 4.93-5.00 (m, 3H); 6.09 (d, 1H, $J = 6.8$); 7.25-7.39 (m, 5H); 8.48 (bs, 1H); 8.65 (s, 1H); 8.89 (s, 1H); 9.51 (bs, 1H); 10.38 (bs, 1H). Anal. ($\text{C}_{21}\text{H}_{25}\text{N}_7\text{O}_5$) Calc'd.: C, 53.50; H, 5.34, N, 20.80. Found: C, 53.55; H, 5.38, N, 20.85.

Example 28: Preparation of N^6 -((S)- α -phenylethylcarbamoyl)-adenosine-5'-N-ethyluronamide

Following the procedure outlined in Example 18, the title compound was prepared in a 72% yield as a white solid, mp 151°C. IR (KBr) cm^{-1} 3550-3100, 1725, 1630, 1525, 1240; ^1H NMR (DMSO d_6) δ : 1.06 (t, 3H, $J = 7$); 1.49 (d, 3H, $J = 8$); 3.15-3.21 (m, 2H); 3.28-3.39 (m, 1H); 4.00 (bs, 1H); 4.35 (s, 1H); 4.61-4.66 (m, 1H); 4.95-5.01 (m, 1H); 6.07 (d, 1H, $J = 6.8$); 7.24-7.39 (m, 5H); 8.48 (bs, 1H), 8.64 (s, 1H); 8.83 (s, 1H); 9.57 (bs, 1H); 10.12 (bs, 1H). Anal. ($\text{C}_{21}\text{H}_{25}\text{N}_7\text{O}_5$) Calc'd.: C, 53.50; H, 5.34, N, 20.80. Found: C, 53.44; H, 5.34, N, 20.77.

Example 29: Preparation of N⁶-(5-methyl-isoxazol-3-yl-carbamoyl)-adenosine-5'-N-ethyluronamide

5 Following the procedure outlined in Example 18, the title compound was prepared in a 58% yield as a white solid, mp 197°C. IR (KBr) cm⁻¹ 3550-3050, 1715, 1620, 1535, 1240; ¹H NMR (DMSO d₆) δ: 1.06 (t, 3H, J = 7); 2.40 (s, 3H); 3.12-3.25 (m, 2H); 4.21-4.23 (m, 1H); 4.36 (d, 1H, J = 2); 4.64-4.70 (m, 1H); 5.68 (m, 2H); 6.09 (d, 1H, J = 6.8); 6.67 (s, 1H); 8.46 (t, 1H, J = 6), 8.74 (s, 1H); 8.84 (s, 1H); 10.75 (bs, 1H); 12.19 (bs, 1H). Anal. 10 (C₁₇H₂₀N₈O₆) Calc'd.: C, 47.22; H, 4.66, N, 25.91. Found: C, 47.14; H, 4.69, N, 25.96.

Example 30: Preparation of N⁶-(1,3,4-thiadiazol-2-yl-carbamoyl)-adenosine-5'-N-ethyluronamide

15 Following the procedure outlined in Example 18, the title compound was prepared in a 65% yield as a white solid, mp 197°C. IR (KBr) cm⁻¹ 3450-3100, 1720, 1635, 1515, 1240; ¹H NMR (DMSO d₆) δ: 1.06 (t, 3H, J = 7); 3.15-3.22 (m, 2H); 3.72 (bs, 2H); 4.21-4.24 (m, 1H); 4.36 (d, 1H, J = 2); 20 4.65-4.70 (m, 1H); 6.12 (d, 1H, J = 6.8); 8.45 (t, 1H, J = 7); 8.80 (s, 1H); 8.89 (s, 1H); 9.19 (s, 1H); 10.35 (bs, 1H); 11.38 (bs, 1H). Anal. (C₁₅H₁₇N₄O₅S) Calc'd.: C, 49.31; H, 4.69, N, 15.33. Found: C, 49.22; H, 4.72, N, 15.33.

25 **Example 31: Preparation of N⁶-(4-n-propoxy-phenylcarbamoyl)-adenosine-5'-N-ethyluronamide**

 Following the procedure outlined in Example 18, the title compound was prepared in a 66% yield as a white solid, mp 270°C. IR (KBr) cm⁻¹ 30 3450-3050, 1720, 1630, 1535, 1240; ¹H NMR (DMSO d₆) δ: 0.95 (t, 3H, J = 7); 1.09 (t, 3H, J = 7); 1.65-1.75 (m, 2H); 3.06-3.23 (m, 2H); 3.67 (bs, 2H); 3.89 (t, 2H, J = 7); 4.18-4.20 (m, 1H); 4.33 (s, 1H); 4.51-4.61 (m, 1H); 6.05 (d, 1H, J = 6.8); 6.91 (d, 2H, J = 9); 7.52 (d, 2H, J = 9); 7.92 (t, 1H, J = 7);

8.22 (s, 1H); 8.52 (s, 1H); 9.65 (bs, 1H); 11.52 (bs, 1H). Anal.
(C₂₂H₂₇N₇O₆) Calc'd.: C, 5.61; H, 5.61, N, 20.19. Found: C, 54.50; H,
5.66, N, 20.06.

5 **Example 32: Preparation of N⁶-bis-(4-nitrophenylcarbamoyl)-
adenosine-5'-N-ethyluronamide**

Following the procedure outlined in Example 18, the title compound
was prepared in a 52% yield as a white solid, mp 213 °C. IR (KBr) cm⁻¹
10 3500-3050, 1720, 1630, 1545, 1340, 1220; ¹H NMR (DMSO d₆) δ: 1.07 (t,
3H, J = 7); 3.12-3.23 (m, 2H); 4.21-4.23 (m, 1H); 4.35 (d, 1H, J = 2); 4.64-
4.69 (m, 1H); 5.69 (bs, 2H); 6.09 (d, 1H, J = 6.8); 7.68 (d, 2H, J = 9); 7.91
(d, 2H, J = 9); 8.18-8.29 (m, 4H); 8.83 (s, 1H); 9.69 (s, 1H); 10.68 (bs, 1H);
12.26 (bs, 1H). Anal. (C₂₆H₂₉N₁₀O₁₀) Calc'd.: C, 48.67; H, 4.56, N, 21.83.
15 Found: C, 48.74; H, 4.60, N, 21.88.

**Example 33: Preparation of N⁶-bis-(5-chloro-pyridin-2-yl-carbamoyl)-
adenosine-5'-N-ethyluronamide**

Following the procedure outlined in Example 18, the title compound
was prepared in a 68% yield as a white solid, mp 220 °C. IR (KBr) cm⁻¹
20 3550-3100, 1725, 1630, 1525, 1250; ¹H NMR (DMSO d₆) δ: 1.06 (t, 3H, J =
7); 3.15-3.22 (m, 2H); 4.22-4.24 (m, 1H); 4.36 (d, 2H, J = 2); 4.64-4.69 (m,
1H); 5.95 (bs, 2H); 6.10 (d, 1H, J = 6.8); 7.80-8.06 (m, 6H); 8.39 (t, 1H, J =
25 7); 8.76 (s, 1H); 8.93 (s, 1H); 10.39 (bs, 1H); 12.11 (bs, 1H). Anal.
(C₂₄H₂₂N₈O₆Cl₂) Calc'd.: C, 48.91; H, 3.76, N, 19.01. Found: C, 49.01;
H, 3.77, N, 18.89.

Example 34. Binding Assays

30

Methods for Performing Receptor Binding Assays

Procedures for preparing rat brain membranes and Chinese hamster ovary (CHO) cell membranes were as reported^{30, 38, 44}. For binding experiments, membrane homogenates were frozen and stored at -20°C for greater than 2 months. Adenosine deaminase (ADA) was purchased from Boehringer Mannheim (Indianapolis, IN). [³H]R-PIA was purchased from Amersham (Arlington Heights, IL), and [³H]CGS 21680 was purchased from DuPont NEN (Boston, MA). [¹²⁵I]-AB-MECA was prepared as described by Olah et al.³⁰

The binding of [¹²⁵I]-AB-MECA to CHO cells stably transfected with the rat A₃ receptor clone or to HEK-293 cells stably expressing the human A₃ receptor, was performed essentially as described^{28, 30, 38, 44}. Assays were performed in 50 mM Tris/10 mM MgCl₂/1 mM EDTA buffer (adjusted to pH 8.26 at 5°C) in glass tubes containing 100 µl of the membrane suspension, 50 µl of [¹²⁵I]-AB-MECA (final concentration 0.3 nM), and 50 µl of inhibitor. Inhibitors were routinely dissolved in DMSO. Concentrations of DMSO in incubations never exceeded 1%; this concentration did not influence [¹²⁵I]-AB-MECA binding. Incubations were carried out in duplicate for 1 h at 37°C, and were terminated by rapid filtration over Whatman GF/B filters using a Brandell cell harvester (Brandell, Gaithersburg, MD). Tubes were washed three times with 3 mL of buffer. Radioactivity was determined in a Beckman gamma 5500B γ-counter.

Nonspecific binding was determined in the presence of 200 µM NECA. K_i values were calculated according to Cheng-Prusoff⁴⁵ assuming a K_d for [¹²⁵I]-AB-MECA of 1.48 nM. Binding of [³H]R-PIA to A₁ receptors from rat cortical membranes and of [³H]CGS 21680 to A_{2A} receptors for rat striatal membranes was performed as described previously.⁴⁶ Adenosine deaminase (2 units/mL) was present during the preparation of the membranes. Additional deaminase was not added during incubation with the radioligand.

Binding of [³⁵S]GTP-γ-S

The binding for [^{35}S]GTP- γ -S (Ameersham, Chicago IL, specific activity 1275 Ci/mmol) was carried out using rat RBL-2H3 mast cell membranes by the general method of Lorenzen et al.⁴⁷ Membranes were suspended in a buffer containing 50 mM Tris, 3 units/mL adenosine deaminase, 100 mM NaCl, and 10 mM MgCl_2 (pH 7.4) and a protein concentration of 1 to 10 μg per tube. The membrane suspension was preincubated with 10 GDP, R-PIA and/or riboflavin in a final volume of 125 μL buffer at 30°C for 60 min and then transferred to ice for 20 min. [^{35}S]GTP- γ -S was added to a final concentration of 0.1 nM in a total volume of 500 μL and the mixture was incubated for 30 min at 30°C. Non-specific binding was determined in the presence of 10 μM GTP- γ -S (Sigma, St. Louis MO). Incubation of the reaction mixture was terminated by filtration over a GF/B glass filter using a Brandell cell harvester and washed with the same buffer.


15

Results and Discussion

20

The derivatives 3a-o (the compounds in Examples 19-33, sequentially) were tested in radioligand binding assays for affinity at rat brain A_1 , $\text{A}_{2\text{A}}$ and A_3 receptors, and the results are summarized in Table 1.

Table 1. Affinities of N⁶-Substituted-carbamoyl-adenosine-5'-uronamide Derivatives in Radioligand Binding Assays at Rat Brain A₁, A_{2A} and A₃ Adenosine Receptors



3a-o

Compound	R	R'	K _i (nM) or % inhibition				A ₁ /A ₃	A _{2A} /A ₃
			K _i (A ₁) ^a	K _i (A _{2A}) ^b	K _i (A ₃) ^c			
IB-MECA	-	-	54 ± 5	56 ± 8	1.1 ± 0.3		49	51
1								
3a	H	4-Biphenyl	5.92 ± 1.04	3580 ± 850	979 ± 22		0.006	3.65

15

Compound	R	R'	K _i (nM) or % inhibition			
			K _i (A ₁) ^a	K _i (A _{2A}) ^b	K _i (A ₃) ^c	A ₁ /A ₃ A _{2A} /A ₃
IB-MECA	-	-	54 ± 5	56 ± 8	1.1 ± 0.3	49 51
1						
3b	H	2,4-Cl-Ph-CH ₂	16 ± 3.3	26.6 ± 8.8	167 ± 7	0.095 0.16
3c	H	4-CH ₃ O-Ph	86.4 ± 14.2	237 ± 46	837 ± 142	0.10 0.28
3d	H	2-Cl-Ph	1170 ± 220	1960 ± 300	754 ± 132	1.55 2.59
3e	H	Ph	252 ± 52	670 ± 211	824 ± 126	0.30 0.81
3f	H	Ph-CH ₂ -NH	171 ± 15	1793 ± 230	38.3 ± 6.4	4.46 46.8
3g	H	4-SO ₂ NH ₂ -Ph-NH	453 ± 141	1180 ± 360	9.73 ± 0.75	46.5 121.2
3h	H	4-CH ₃ CO-Ph-NH	72.7 ± 11.8	1050 ± 270	20.9 ± 7.51	3.47 50.2
3i	H	(R)-α-Phenylethyl-NH	433 ± 171	279 ± 171	16.3 ± 3.7	26.5 17.1
3j	H	(S)-α-Phenylethyl-NH	537 ± 33	2970 ± 930	319 ± 134	1.68 9.3
3k	H	5-Me-Isoxazol-3-yl-NH	146 ± 39	884 ± 232	532 ± 240	0.27 1.66
3l	H	1,3,4-Thiadiazol-2-yl-NH	208 ± 42	917 ± 254	5550 ± 2800	0.037 0.16
3m	H	4-nC ₃ H ₇ O-Ph-NH	257 ± 43	255 ± 85	107 ± 34	2.3 2.4
3n	4-NO ₂ -Ph-NH-CO	4-NO ₂ -Ph-NH	89.1 ± 8.8	2530 ± 200	168 ± 42	0.53 15

5

38

15

Compound	R	R'	K _i (nM) or % inhibition				A ₁ /A ₃	A _{2A} /A ₃
			K _i (A ₁) ^a	K _i (A _{2A}) ^b	K _i (A ₃) ^c			
IB-MECA	-	-	54 ± 5	56 ± 8	1.1 ± 0.3	49	51	
1								
3o	5-Cl-	5-Cl-Pyridin-2-yl-NH	761 ± 187	799 ± 191	5700 ± 100	0.13	0.14	
	Pyridin-2-yl-	NH-CO						

3o

- a Displacement of specific [³H]R-PIA binding (A₁) in rat brain membranes, expressed as K_i ± SEM in nM (n=3-6).
- b Displacement of specific [³H]CGS 21680 binding (A_{2A}) in rat striatal membranes expressed as K_i ± SEM in nM (n=3-6).
- c Displacement of specific [¹²⁵I]AB-MECA binding at rat A₃ receptors expressed in CHO cells, expressed as K_i ± SEM in nM (n=3-6).

5

Baraldi demonstrated that certain N⁶-substituted-arylcarbamoyl adenosine-uronamides are full agonists in inhibiting adenylate cyclase via rat A₃ receptors, but that the compounds showed relatively high activity with respect to adenosine A₁ receptors³⁹. There is a similarity between A₁ and A₃ receptors. Small modifications in the length of chain at the N⁶ position can alter the affinity versus A₁ or A₃ receptors. The present binding data demonstrate that the presence of aryl- or alkaryl-carbamoylic chains at the N⁶-position bring about an interaction with the A₃ receptor subtype. Also, the data demonstrate that heteroaryl-carbamoyl, heteroaryl-carboxamide, aryl-carboxamide, and alkaryl-carboxamide substitutions at the N⁶-position bring about an interaction with the A₁ receptor subtype.

In the series of urea-derivatives (Compounds 3f-o) a substituted-phenyl (Compounds 3g, h and m) or substituted-benzyl (Compounds 3f, i and j) group leads to relatively high affinity and selectivity at A₃ adenosine receptors as compared with the affinity for the A₁ and A₂ receptors.

Derivatives 3g, 3h and 3m showed a relatively high affinity (9.7-107 nM) at A₃ receptors with varying degrees of A₃/A₁ selectivity. In particular, compound 3g was less active than IB-MECA (9.7 nM vs. 1.1 nM) but showed selectivity for A₃ vs. either A₁ or A_{2A} receptors which is comparable to IB-MECA.

Compound 3g also showed a relatively high affinity, in the nanomolar range, at human A₃ adenosine receptors (56.1 ± 9.1 nM), confirming the relatively high affinity of this compound versus this receptor subtype, independent of species. The lipophilicity of para substituents on a phenyl ring play a significant role in A₃ affinity.

Substitution of the phenyl ring with a heterocycle (as in Compounds 3k-l) causes the compounds to lose affinity and selectivity for A₃ receptors, and increase in affinity for the A₁ receptor subtype.

5 Disubstitution at the N⁶ position (for example, Compounds 3n-o) have diminished activity at A_{2A} and A₃ receptors, but have relatively high affinity for the A₁ receptor.

10 In the benzylic series, Compounds 3f, i, and j showed an SAR pattern comparable to the phenylic series in Baraldi³⁹. In fact, the unsubstituted Compound 3f showed affinity and selectivity at A₃ receptor very similar to the previously reported phenyl derivative.³⁹

15 The stereochemistry of substituents at the N⁶ position is also important in determining the affinity of these compounds for the A₃ receptor. For example, comparing the two diastereomeric Compounds 3i and 3j in which substituents at N⁶ are of opposite configuration, the R-isomer (3i) is more potent and selective than the S-isomer (3j) at the rat A₃ adenosine receptor subtype. Those of skill in the art can separate compounds with
20 stereocenters adjacent to the N⁶ position using routine enantiopurification methods and evaluate the individual stereoisomers for their affinity and selectivity using no more than routine experimentation.

25 A functional assay indicated that Compound 3g acted as a full agonist at rat A₃ receptor. The assay involved the agonist-induced inhibition of binding of guanine nucleotide to rat RBL-2H3 mast cell membranes, which contain a high density of A₃ receptors.

30 Figure 1 shows that compound 3g was about as effective as the potent A₃ agonists I-AB-MECA and Cl-IB-MECA increased binding of [³⁵S]GTP-γ-S in a dose dependent manner and with greater potency than NECA.

Example 35. Pharmaceutical Formulations**(A) Transdermal System - for 1000 patches**

5

Ingredients	Amount
Active compound	100g
Silicone fluid	450g
Colloidal silicon dioxide	2g

10

The silicone fluid and active compound are mixed together and the colloidal silicone dioxide is added to increase viscosity. The material is then dosed into a subsequent heat sealed polymeric laminate including the following: polyester release liner, skin contact adhesive composed of silicone or acrylic polymers, a control membrane which is a polyolefin, and an impermeable backing membrane made of a polyester multilaminate. The resulting laminated sheet is then cut into 10 sq. cm patches

15

(B) Oral Tablet - For 1000 Tablets

20

Ingredients	Amount
Active compound	50g
Starch	50g
Magnesium Stearate	5g

25

The active compound and the starch are granulated with water and dried. Magnesium stearate is added to the dried granules and the mixture is thoroughly blended. The blended mixture is compressed into tablets.

30

(C) Injection - for 1000, 1mL Ampules

Ingredients	Amount
Active compound	10g
Buffering Agents	q.s.
Propylene glycol	400mg
Water for injection	q.s.1000mL

The active compound and buffering agents are dissolved in the propylene glycol at about 50°C. The water for injection is then added with stirring and the resulting solution is filtered, filled into ampules, sealed and sterilized by autoclaving.

(D) Continuous Injection - for 1000 mL

Ingredients	Amount
Active compound	10g
Buffering agents	q.s.
Water for injection	q.s.1000mL

REFERENCES:

The following references have been referred to herein. These references are incorporated herein by reference in their entirety.

1. Libert, F., Schiffmann, S.N.; Lefort, A., Parmentier, M; Gerard, C.; Dumont, J E.; Vanderhaeghen, J.J.; Vassart, G. The orphan receptor

cDNA RDC7 encodes an A₁ adenosine receptor. EMBO J. 1991, 10, 1677-1682.

2. Maenhaut, C., Sande, J.V.; Libert, F; Abramowicz, M.;
5 Parmentier, M.; Vanderhaeghen, J.J.; Dumont, J.E.; Vassart, G.; Schiffmann, S. RDC8 codes for an adenosine A₂ receptor with physiological constitutive activity. Biochem. Biophys. Res. Commun. 1990, 173, 1169-1178.
3. Stehle, J.H.; Rivkees, S.A.; Lee, J.J.; Weaver, D.R.; Deeds, J.D.;
10 Reppert, S.M. Molecular cloning and expression of the cDNA for a novel A₂-adenosine receptor subtype. Mol. Endocrinol. 1992, 6, 384-393.
4. Zhou, Q.Y.; Li, C.Y.; Olah, M.E.; Johnson, R.A.; Stiles, G.L.;
Civelli, O. Molecular cloning and characterization of an adenosine receptor-
15 the A₃ adenosine receptor. Proc. Natl. Acad. Sci. USA 1992, 89, 7432-7436.
5. Londos, C., Cooper, D.M.; Wolff, J. Subclasses of external
adenosine receptors. Proc. Natl. Acad. Sci. USA 1980, 77, 2551-2554.
- 20 6. Hamprecht, B.; Van Calker, D. Nomenclature of adenosine
receptors. Trends Pharmacol. Sci. 1985, 6, 153-154.
7. Ukena, D.; Olsson, R.A.; Daly, J.W. Definition of subclasses of
adenosine receptors associated with adenylate cyclase: interaction of
25 adenosine analogs with inhibitory A₁ receptors and stimulatory A₂ receptors.
Can. J. Physiol. Pharmacol. 1987, 65, 365-376.
8. Ramkumar, V., Stiles, G.L.; Beaven, M.A.; Ali, H. The A₃AR is
the unique adenosine receptor which facilitates release of allergic mediators
30 in mast cells. J Biol. Chem. 1993, 268, 6871-6890.

9. Abbracchio, M.P.; Brambilla, R.; Ceruti, S.; Kim, H.O.; von Lubitz, D.K.J.E.; Jacobson, K.A., Cattabeni, F. G-protein-dependent activation of phospholipase C by adenosine A₃ receptors in rat brain. Mol. Pharmacol. 1995, 48, 1038-1045.

5

10. Bruns, R.F.; Daly, J.W.; Snyder, S.H. Adenosine receptors in brain membranes: binding of N-cyclohexyl [³H] adenosine and 1,3-diethyl-8(³H) phenylxanthine. Proc. Natl. Acad. Sci. USA 1980, 77, 5547-5551.

10

11. Olsson, R.A.; Kusachi, S.; Thompson, S.D.; Ukena, D., Padgett, W.; Daly, J.W. N⁶-substituted N-alkyladenosine-5¹-uronamides: bifunctional ligands having recognition groups for A₁ and A₂ adenosine receptors. J. Med. Chem. 1986, 29, 1683-1689.

15

12. Jacobson, K.A.; Nikodijevic, O.; Ji, X.D.; Berkich, D.A.; Eveleth, D.; Dean, R.L.; Hiramatsu, K.; Kassel, N.F.; van Galen, P.J.M.; Lee, K.S.; Bartus, R.T.; Daly, J.W.; Lanoue, K.F.; Maillard, M. Synthesis and biological activity of N⁶-(p-sulfophenyl) alkyl and N⁶-p-sulfoalkyl derivatives of adenosine. Water soluble and peripherally selective adenosine agonists. J. Med. Chem. 1992, 35, 4143-4149.

20

13. Shamim, M.T.; Ukena, D.; Padgett, W.L.; Hong, O.; Daly, J.W. 8-Aryl-and 8cycloalkyl-1,3-dipropylxanthines: further potent and selective antagonists for A₁-adenosine receptors. J. Med. Chem. 1988, 31, 613-617.

25

14. Linden, J.; Patel, A.; Earl, C.Q.; Craig, R.H.; Daluge, S.M. (¹²⁵I)-labeled 8-phenylxanthine derivatives: antagonist radioligands for adenosine A₁ receptors. J. Med. Chem. 1988, 31, 745-751.

30

15. Jacobson, K.A. de la Cruz., R.; Schulick, R.; Kiriasis, K.L.; Padgett, W.; Pfeleiderer, W.; Kirk, K.L.; Neumeyer, J.L., Daly, J.W. 8-

substituted xanthines as antagonists at A₁ and A₂ adenosine receptors.
Biochem. Pharmacol. 1988, 37, 3653-3661.

5 16. Cristalli, G.; Eleuteri, A.; Vittori, S.; Volpini, R.; Lohse, M.J.;
Klotz, K.N. 2-alkynyl derivatives of adenosine and adenosine-5'-N-
ethyluronamide as selective agonists at A₂ adenosine receptors. J. Med.
Chem. 1992, 35, 2363-2368.

10 17. Jarvis, M.F.; Schulz, R.; Hutchison, A.J.; Do, U.H.; Sills, M.A.;
Williams, M. [³H]CGS 21680, a selective A₂ adenosine receptor agonist
directly labels A₂ receptors in rat brain. J. Pharmacol. Exp. Ther. 1989, 251 ,
888-893.

15 18. Shimada, J.; Suzuki, K.; Nonaka, H.; [shii, A.; [chikawa, S. (E)-
1,3-dialkyl-7-methyl-8-(3,4,5'-trimethoxystyryl)xanthines: potent and
selective adenosine A₂ antagonists. J. Med. Chem. 1992, 35, 2342-2345.

20 19. a) Jacobson, K.A.; Gallo-Rodriguez, C.; Melman, N.; Fischer,
B.; Malillard, M.; van Bergen, A.; van Galen, P.J.M.; Kar-ton, Y. Structure-
activity relationship of 8-styrylxanthines as A₂-selective adenosine
antagonists. J. Med. Chem. 1993, 36, 1333-1342; b) Baraldi, P.G.;
Manfredini, S.; Simoni, D.; Zappaterra, L.; Zocchi, C.; Dionisotti, S.;
Ongini, E. Synthesis and activity of new pyrazolo[4,3-e]- 1,2,4-triazolo[1,5-
c] pyrimidine and 1,2,3-triazolo[4,5-e] 1,2,4-triazolo [1,5-c]pyrimidine
25 displaying potent and selective activity as A_{2A} adenosine receptor
antagonist. Bioorg. Med. Chem. Lett. 1994, 4, 2539-2544; c) Baraldi, P.G.;
Cacciari, B.; Spalluto, G.; Borioni, A.; Vizio, M.; Dionisotti, S.; Ongini, E.
Current developments of A_{2A} adenosine receptor antagonists- Current
Medicinal Chemistry 1995, 2, 707-722.

30

20. Ferré, S.; O'Connor, W.T.; Snaprud, P.; Ungerstedt, U.; Fuxe, K. Antagonistic interactions between adenosine A_{2A} receptors and dopamine D₂ receptors in the ventral striopallidal system-implications for the treatment of schizophrenia. Neurosci. 1994, 63, 765-773.

5

21. Schingnitz, G.; Kufner-Mühl, U.; Ensinger, H.; Lehr, E.; Kuhn, F.J. Selective A₁- antagonists for treatment of cognitive deficits. Nucleosides Nucleotides 1991, 10, 1067-1076.

10

22. Bridges, A.J.; Moos, W.H.; Szotek, D.L.; Trivedi, B.K.; Bristol, J.A.; Heffner, T.G.; Bruns, R.F.; Downs, D.A. N⁶-(2,2-diphenylethyl)adenosine, a novel adenosine receptor agonist with anti psychotic-like activity. J. Med. Chem. 1987, 30, 1709-1711.

15

23. Jacobson, K.A.; Trivedi, B.K.; Churchill, P.C.; Williams, M. Novel therapeutics acting via purine receptors. Biochem. Pharmacol. 1991, 41, 1399- 1410.

20

24. Jacobson, K.A.; Kim, H.O.; Siddiqi, S.M.; Olah, M.E.; Stiles, G.; von. Lubitz, D.K.J. A₃ adenosine receptors: design of selective ligands and therapeutic prospects. Drugs Future 1995, 20, 689-699.

25

25. Ji, X.D.; Melman, N.; Jacobson, K.A. Interactions of flavonoids and other phytochemicals with adenosine receptors. J. Med. Chem. 1996, 39, 781-788.

30

26. Karton, Y.; Jiang, J.-I.; Ji, X.D.; Melman, N.; Olah, M.E.; Stiles, G.L.; Jacobson, K.A. Synthesis and biological activities of flavonoid derivatives as A₃ adenosine receptor antagonists. J. Med. Chem. 1996, 39, 2293-2301.

27. van Rhee, A.M.; Jiang, J.-I.; Melman, N.; Olah, M.E.; Stiles, G.L.; Jacobson, K.A. Interaction of 1,4-dihydropyridine and pyridine derivatives with adenosine receptors: selectivity for A₃ receptors. J. Med. Chem. 1996, 39, 2980-2989.
28. Kim, Y.-C.; Ji, X.D.; Jacobson, K.A. Derivatives of the triazoloquinazoline adenosine antagonist (CGS 15943) are selective for the human A₃ receptor subtype. J. Med. Chem. 1996, 39, 4142-4148.
29. Kim, H.O.; Ji, X.D.; Siddiqi, S.M.; Olah, M.E.; Stiles, G.L.; Jacobson, K.A. 2-substitution of N⁶-benzyladenosine-5'-uronamides enhances selectivity for A₃ adenosine receptors. J. Med. Chem. 1994, 37, 3614-3621.
30. Olah, M.E.; Gallo-Rodriguez, C.; Jacobson, K.A.; Stiles, G.L. [¹²⁵I]AB-MECA, a high affinity radioligand for the rat A₃ adenosine receptor. Mol. Pharmacol. 1994, 45, 978-982.
31. Linden, J. Cloned adenosine A₃ receptors-pharmacological properties, species, differences and receptor functions. Trends Pharmacol. Sci. 1994, 15, 298-300.
32. Hannon, J.P.; Pfannkuche, H.J.; Fozard, J.R. A role for mast cells in adenosine A₃ receptor-mediated hypotension in the rat. Br. J. Pharmacol. 1995, 115, 945-952.
33. Fozard, J.R.; Pfannkuche, H.J.; Schuurmari, H.J. Mast cell degranulation following adenosine A₃ receptor activation in rats. Eur. J. Pharmacol. 1996, 298, 293-297.


34. Jacobson, K.A.; Nikodijevic, O.; Shi, D.; Gallo-Rodriguez, C.; Olah, M.E.; Stiles, G.L.; Daly, J.W. A role for central A₃-adenosine receptors: Mediation of behavioral depressant effects. FEBS Lett 1993, 336, 57-60.
- 5 35. von Lubitz, D.K.J.E.; Lin, R.C.S.; Popik, P.; Carter, M.F.; Jacobson, K.A. Adenosine A₃ receptor stimulation and cerebral ischemia Eur. J. Pharmacol. 1994, 263,59-67.
- 10 36. Kohno, Y.; Sei, Y.; Koshiba, M.; Kim, H.O.; Jacobson, K.A. Induction of apoptosis in HL-60 human promyelocytic leukemia cells by selective adenosine A₃ receptor agonists. Biochem. Biophys. Res. Commun. 1996, 219, 904-910.
- 15 37. Jacobson, K.A.; van Galen, P.J.M.; Williams, M. Perspective Adenosine Receptors: pharmacology, structure-activity relationship and therapeutic potential. J. Med. Chem. 1992,35, 407-422.
- 20 38. Gallo- Rodriguez, C.; Ji, X.D.; Melman, N.; Siegman, B.D.; Sanders, L.H.; Orlina, J.; Pu, Q.L.; Olah, M.E.; van Galen, P.J.M.; Stiles, G.L.; Jacobson, K.A. Structure-activity relationship of N⁶-benzyladenosine-5'-uronamides as A₃-selective adenosine agonists. J. Med. Chem. 1994, 37, 636-646.
- 25 39. Baraldi, P.G.; Cacciari, B.; Spalluto, G.; Ji, X.D.; Olah, M.E.; Stiles, G.; Dionisotti, S.; Zocchi, S.; Ongini, E.; Jacobson, K.A. Novel N⁶-(substituted-phenylcarbamoyl)adenosine-5'-uronamides as potent agonists for A₃ adenosine receptors. J. Med. Chem. 1996, 39, 802-806.

40. Kurita, K.; Iwak-ura, Y. Trichloromethyl -Chloroformate as a phosgene equivalent: 3 -isocyanatopropanoylchloride. Org. Synth. Coll. vol. VI, 1988, 715-718.
- 5 41. Siddiqi, S.M.; Pearlstein, R.A.; Sanders, L.H.; Jacobson, K.A. Comparative molecular field analysis of selective A₃ adenosine receptor agonists. Bioorg. Med. Chem. 1995, 3, 1331-1343.
- 10 42. van Galen, P.J.M.; Stiles, G.L.; Michaels, G.; Jacobson, K.A. Adenosine A₁ and A₂ receptors: structure-function relationships. Med. Res. Rev. 1992, 12, 423-471.
43. Muller, C.E.; Scior, T. Adenosine receptors and their modulators. Pharmaceutica Acta Helveticae 1993,68, 77-111.
-
- 15 44. van Galen, P.J.M.; van Bergen, A.H.; Gallo-Rodriguez, C.; Melman, N.; Olah, M.E.; Ijzerman, A.P.; Stiles, G.L.; Jacobson, K.A. A binding site model and structure-activity relationship for the rat A₃ adenosine receptor. Mol. Pharmacol. 1994, 45, 1101-1111.
- 20 45. Cheng, Y.C.; Prusoff, H.R. Relationships between the inhibition constant (K_i) and the concentration of inhibitor which causes 50% inhibition (IC₅₀) of an enzymatic reaction. Biochem. Pharmacol. 1973, 22, 3099-3108.
- 25 46. van Bergen, A.; van Galen, P.J.M., Stiles, G.L.; Jacobson, K.A. A₃ receptors: Structure activity relationship and molecular modeling. ACS 206th National Meeting. Chicago, IL, August 1993, Abstract MEDI217.
- 30 47. Lorenzen, A.; Guerra, L.; Vogt, H.; Schwabe, U. Interaction of full and partial agonists of the A₁ adenosine receptor with receptor/G protein complexes in rat brain membranes. Mol. Pharmacol. 1996, 49, 915-926.

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific
5 embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed is:

1. A compound of the following formula:



wherein:

R is hydrogen, alkyl, substituted alkyl, alkaryl or aryl;

R¹ is heteroaryl-NR-C(X)-, heteroaryl-C(X)-, alkaryl-NR-C(X)-, substituted alkaryl-NR-C(X)-, aryl-NR-C(X)-, aryl-C(X)-, alkaryl-C(X)-, or substituted alkaryl-C(X)-,

R² is hydrogen, alkyl, substituted alkyl, alkaryl or aryl-NH-C(X)-,

R³ and R⁴ are independently selected from the group consisting of hydroxy, hydrogen, halo, ether, ester, azide, alkyl, alkoxy, carboxy, nitrile, nitro, trifluoro, aryl, alkaryl, thio, thioester, thioether, amine and amide, and

X is O, S, or NR, with the proviso that, when R² is H, R¹ is aryl-NH-C(X)- and X is O, the substituents on the aryl ring are not halo, methoxy or trifluoromethyl.

2. The compound of claim 1 wherein R is hydrogen.
3. The compound of claim 1 wherein R¹ is alkaryl-NR-C(X)-.
4. The compound of claim 1 wherein R¹ is substituted alkaryl-NR-C(X)-.
5. The compound of claim 1 wherein R¹ is aryl-NR-C(X)-.
6. The compound of claim 1 wherein R¹ is aryl-C(X)-.
7. The compound of claim 1 wherein R¹ is alkaryl-C(X)-.
8. The compound of claim 1 wherein R¹ is substituted alkaryl-C(X)-.
9. The compound of claim 1 wherein R¹ is heteroaryl-NR-C(X)-.
10. The compound of claim 1 wherein R¹ is heteroaryl-C(X)-.
11. The compound of claim 1 wherein R² is hydrogen.
12. The compound of claim 1 wherein X is O.
13. A compound selected from the group consisting of:
 - N⁶-(4-biphenyl-carbonylamino)-adenosine-5'-N-ethyluronamide,
 - N⁶-(2,4-dichlorobenzyl-carbonylamino)-adenosine-5'-N-ethyluronamide,
 - N⁶-(4-methoxyphenyl-carbonylamino)-adenosine-5'-N-ethyluronamide,
 - N⁶-(4-chlorophenyl-carbonylamino)-adenosine-5'-N-ethyluronamide,
 - N⁶-(phenyl-carbonylamino)-adenosine-5'-N-ethyluronamide,
 - N⁶-(benzylcarbamoylamino)-adenosine-5'-N-ethyluronamide,
 - N⁶-(4-sulfonamido-phenylcarbamoyl)-adenosine-5'-N-ethyluronamide,
 - N⁶-(4-acetyl-phenylcarbamoyl)-adenosine-5'-N-ethyluronamide,
 - N⁶-((R)- α -phenylethylcarbamoyl)-adenosine-5'-N-ethyluronamide,
 - N⁶-((S)- α -phenylethylcarbamoyl)-adenosine-5'-N-ethyluronamide,
 - N⁶-(5-methyl-isoxazol-3-yl-carbamoyl)-adenosine-5'-N-ethyluronamide,
 - N⁶-(1,3,4-thiadiazol-2-yl-carbamoyl)-adenosine-5'-N-ethyluronamide,
 - N⁶-(4-n-propoxy-phenylcarbamoyl)-adenosine-5'-N-ethyluronamide,

N⁶-bis-(4-nitrophenylcarbamoyl)-adenosine-5'-N-ethyluronamide,
and

N⁶-bis-(5-chloro-pyridin-2-yl-carbamoyl)-adenosine-5'-N-ethyluronamide.

14. A method of treating tumors, hypertension, mast cell degranulation, or cardiac hypoxia, or providing protection against cerebral ischemia, comprising administering to a patient in need of treatment thereof an effective amount of a compound of claim 1 which is an agonist or partial agonist of the A₃ receptor.

15. The method of claim 14 wherein R is hydrogen.

16. The method of claim 14 wherein R¹ is alkaryl-NR-C(X)-.

17. The method of claim 14 wherein R¹ is substituted alkaryl-NR-C(X)-.

18. The method of claim 14 wherein R¹ is aryl-NR-C(X)-.

19. The method of claim 14 wherein R² is hydrogen.

20. The method of claim 14 wherein X is O.

21. The method of claim 14 wherein the compound is selected from the group consisting of:

N⁶-(benzylcarbamoylamino)-adenosine-5'-N-ethyluronamide,

N⁶-(4-sulfonamido-phenylcarbamoyl)-adenosine-5'-N-ethyluronamide,

N⁶-(4-acetyl-phenylcarbamoyl)-adenosine-5'-N-ethyluronamide,

N⁶-((R)- α -phenylethylcarbamoyl)-adenosine-5'-N-ethyluronamide,

N⁶-((S)- α -phenylethylcarbamoyl)-adenosine-5'-N-ethyluronamide,

N⁶-(4-n-propoxy-phenylcarbamoyl)-adenosine-5'-N-ethyluronamide,

and

N⁶-bis-(4-nitrophenylcarbamoyl)-adenosine-5'-N-ethyluronamide,

22. A method for providing protection against hypoxia, ischemia induced injuries, cardioprotection, neuroprotection, pain management, reduction of free fatty acids, triglycerides, or glucose levels, adjunct therapy

in diabetes, treatment of adenosine-sensitive cardiac dysarrhythmias, treatment of GI disorders, treatment of convulsion, treatment of glaucoma; treatment of sleep apnea; treatment of peroxysmal supraventricular tachycardia, or treatment of inflammation comprising administering to a patient in need of treatment thereof an effective amount to treat the disorder of a compound of Formula I which is active as an agonist or partial agonist of the A₁ receptor.

23. The method of claim 23 wherein R¹ is aryl-C(X)-.

24. The method of claim 23 wherein R¹ is alkaryl-C(X)-.

25. The method of claim 23 wherein R¹ is substituted alkaryl-C(X)-.

26. The method of claim 23 wherein R¹ is heteroaryl-NR-C(X)-.

27. The method of claim 23 wherein R¹ is heteraryl-C(X)-.

28. The method of claim 23 wherein the compound is selected from the group consisting of:

N⁶-(4-biphenyl-carbonylamino)-adenosine-5'-N-ethyluronamide,

N⁶-(2,4-dichlorobenzyl-carbonylamino)-adenosine-5'-N-ethyluronamide,

N⁶-(4-methoxyphenyl-carbonylamino)-adenosine-5'-N-ethyluronamide,

N⁶-(4-chlorophenyl-carbonylamino)-adenosine-5'-N-ethyluronamide,

N⁶-(phenyl-carbonylamino)-adenosine-5'-N-ethyluronamide,

N⁶-(5-methyl-isoxazol-3-yl-carbamoyl)-adenosine-5'-N-ethyluronamide,

N⁶-(1,3,4-thiadiazol-2-yl-carbamoyl)-adenosine-5'-N-ethyluronamide, and

N⁶-bis-(5-chloro-pyridin-2-yl-carbamoyl)-adenosine-5'-N-ethyluronamide.

**N⁶-Substituted-Adenosine-5'-Uronamides
as Adenosine Receptor Modulators**

ABSTRACT

A series of adenosine-5'-uronamide derivatives bearing N⁶-arylurea, alkarylurea, heteroarylurea, arylcarbonyl, alkarylcarbonyl or heteroarylcarbonyl groups which have affinity and, in some cases, selectivity for the adenosine A₁ or A₃ receptors are disclosed. These compounds can be used in a pharmaceutical composition to treat disorders caused by excessive activation of the A₁ or A₃ receptors, or can be used in a diagnostic application to determine the relative binding of other compounds to the A₁ or A₃ receptors.






FIG. 1




5a-e
R = aryl, aralkyl



R = Aryl, Aralkyl; R' = H 5f-m
R = Aryl, R' = R-NH-CO 5n-o


Dioxane/reflux
18h

Dioxane/reflux
18h



+ R-COCl

4





+ R-N=C=O

4

FIG. 2A

3 / 3



a-0

$R = H, Aryl-NH-CO, Aralkyl-NH-CO$
 $R' = Aryl, Aralkyl, Aryl-NH-CO, Aralkyl-NH-CO$

3a-0

FIG.2B

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/16053

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/70

US CL : 536/26.12, 26.13

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/26.12, 26.13

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

USPATFULL, APS, HCAPLUS, REGISTRY

Search terms: uronamide(s), adenosine, adenosine 5' uronamide, antiviral, anticancer

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---- Y	US 5,310,731 A (OLSSON et al.) 05 May 1994, see entire document, especially columns 1-9.	1-13 ----- 14-21
X ---- Y	BARALDI et al. Novel N ⁶ -(Substituted-phenylcarbamoyl)Adenosine-5'-Uronamides as Potent Agonists for A ₃ Adenosine Receptors. Journal of Medicinal Chemistry. February 1996, Vol. 39, No. 3, pages 802-805, especially page 803.	1-13, 22-28 ----- 14-21



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

02 OCTOBER 1998

Date of mailing of the international search report

28 OCT 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

HOWARD OWENS

Telephone No. (703) 308-1234